

INFORMATION  
ANALYSIS  
DIRECTORATE



# Health Inequalities

## Annual Report 2021

A product of the NI Health and Social Care Inequalities Monitoring System



Department of  
**Health**

An Roinn Sláinte  
Mánnystrie O Poustie  
[www.health-ni.gov.uk](http://www.health-ni.gov.uk)

Annual Publication  
April 2021

# Health Inequalities

Annual Report 2021

Authors: Philip Carson, Hannah Blakley & Caolan Laverty

**Public Health Information & Research Branch, Information Analysis Directorate**

**Lead Statistician: Bill Stewart**

Department of Health  
Castle Buildings, Belfast BT4 3SQ

☎ Tel: 02890 522 501

☎ Ext: 22501

✉ Email: [healthinequalities@health-ni.gov.uk](mailto:healthinequalities@health-ni.gov.uk)

<http://www.health-ni.gov.uk/topics/dhssps-statistics-and-research/health-inequalities-statistics>

For information on other Government statistics contact:  
The Northern Ireland Statistics and Research Agency (NISRA)  
Colby House, Stranmillis Court, Belfast, BT9 5RR

☎ Tel: 02890388400

✉ Email: [info@nisra.gov.uk](mailto:info@nisra.gov.uk)

[www.nisra.gov.uk](http://www.nisra.gov.uk)

© Crown Copyright 2021. Published with the permission of the Controller of Her Majesty's Stationary Office.

This publication is Crown copyright and may be reproduced free of charge in any format or medium.  
Any material used must be acknowledged and sourced to the NI Health and Social Care Inequalities Monitoring System.

**Information Analysis Directorate (IAD)** sits within the **Department of Health (DoH)** and carries out various statistical work and research on behalf of the department. It comprises four statistical areas: Hospital Information, Community Information, Public Health Information & Research and Project Support Analysis.

IAD is responsible for compiling, processing, analysing, interpreting and disseminating a wide range of statistics covering health and social care.

The statisticians within IAD are out-posted from the Northern Ireland Statistics & Research Agency (NISRA) and our statistics are produced in accordance with the principles and protocols set out in the UK Code of Practice for Official Statistics.

### About Public Health Information and Research Branch

The role of Public Health Information and Research Branch (PHIRB) is to support public health policy development through managing the public health survey function while also providing analysis and monitoring data. The head of the branch is the Principal Statistician, Mr. Bill Stewart.

In support of the public health survey function, PHIRB is involved in the commissioning, managing and publishing of results from departmental funded surveys, such as the Health Survey Northern Ireland, All Ireland Drug Prevalence Survey, Young Persons Behaviour & Attitudes Survey, Patient Experience Surveys and the Adult Drinking Patterns Survey.

The branch also houses the NI Health and Social Care Inequalities Monitoring System which covers a range of different health inequality/equality based projects conducted for both the region as well as for more localised area levels. In addition, PHIRB is responsible for the production of official life expectancy estimates for NI, and areas within the region.

PHIRB provides support to a range of key DoH NI strategies including Making Life Better, a 10 year cross-departmental public health strategic framework as well as a range of other departmental strategies such as those dealing with suicide, sexual health, breastfeeding, tobacco control and obesity prevention. It also has a key role in supporting the Departmental Alcohol and Drug Strategy, by maintaining and developing key departmental databases such as, the Substance Misuse Database, Impact Measurement Tool and the Census of Drug & Alcohol Treatment Services, which are all used to monitor drug misuse and treatments across Northern Ireland. In addition to Departmental functions, PHIRB also support the executive level Programme for Government and its strategic outcomes through a series of performance indicators.

## CONTENTS

	<b>PAGE NUMBER</b>
KEY FINDINGS	5
INTRODUCTION	7
FORMAT OF THE REPORT	7
REGIONAL HEALTH INEQUALITIES	11
SUMMARY OF CHANGES IN REGIONAL INEQUALITY GAPS	12
LIFE EXPECTANCY & GENERAL HEALTH	14
PREMATURE MORTALITY	17
MAJOR DISEASES	20
HOSPITAL ACTIVITY	22
MENTAL HEALTH	24
ALCOHOL, SMOKING & DRUGS	25
PREGNANCY & EARLY YEARS	28
CHILDHOOD OBESITY	30
SUB-REGIONAL HEALTH INEQUALITIES	32
SUMMARY OF CHANGES IN SUBREGIONAL INEQUALITY GAPS	33
COMPARISON OF SUB-REGIONAL HEALTH OUTCOMES AGAINST REGIONAL AVERAGE	34
LARGEST DEPRIVATION INEQUALITY GAPS IN EACH AREA	35
HEALTH & SOCIAL CARE TRUSTS	
BELFAST	36
NORTHERN	37
SOUTH EASTERN	38
SOUTHERN	39
WESTERN	40
LOCAL GOVERNMENT DISTRICTS	
ANTRIM & NEWTOWN ABBEY	41
ARDS & NORTH DOWN	42
ARMAGH CITY, BANBRIDGE & CRAIGAVON	43
BELFAST LGD	44
CAUSEWAY COAST & GLENS	45
DERRY CITY & STRBANE	46
FERMANAGH & OMAGH	47
LISBURN & CASTLEREAGH	48
MID & EAST ANTRIM	49
MID ULSTER	50
NEWRY, MOURNE & DOWN	51
<b>APPENDICIES</b>	
APPENDIX A: SOCIAL GRADIENT OF HEALTH	52
APPENDIX B: POPULATION ATTRIBUTABLE RISK (PAR) OF DEPRIVATION	58
APPENDIX C: ADDITIONAL INDICATORS	59
APPENDIX D: URBAN RURAL ANALYSIS	61
APPENDIX E: TECHNICAL NOTES & DEFINITIONS	62

## KEY FINDINGS – REGIONAL (NI)

- In 2017-19, male life expectancy at birth has continued to improve in NI and its most and least deprived areas, with no notable change in the deprivation gap (7.0 years) observed since 2013-15. Female life expectancy improved in NI, with no notable change in the most and least deprived areas or the deprivation gap (4.8 years).
- Between 2013-15 and 2017-19 there was no change in male or female healthy life expectancy in NI and its most and least deprived areas, including no notable change in the deprivation gap. Similarly, disability-free life expectancy saw no notable changes across NI, its most and least deprived areas or the deprivation gap for both males and females.
- For indicators of premature mortality, rates generally decreased over the period in NI and its most and least deprived areas. However, large inequality gaps continue to persist, with the rate of respiratory mortality among under 75s in the most deprived areas three and a half times that in the least deprived. Across indicators of premature mortality, the inequality gaps narrowed or remained broadly similar except for potential years of life lost and death rates for preventable causes where the gap widened.
- The inequality gap for crude death rate for intentional self-harm narrowed as a result of positive changes in the most deprived areas, where the rate is approximately double that in the least deprived areas.
- Alcohol and drug related indicators continue to show some of the largest health inequalities monitored in NI, with rates in the most deprived areas five times that in the least deprived for drug related mortality and four times that for alcohol specific mortality.
- In 2019, within the most deprived areas the proportion of births where the mother reported smoking during pregnancy in the most deprived areas was four and a half times the rate in the least deprived areas, with the inequality gap widening over the last five years.

Most Notable Inequality Gaps		Most Notable Narrowing of Gaps	Most Notable Widening of Gaps
Female HLE	15.4 yrs	SAR – Self-Harm	SDR – Drug Misuse
Male HLE	13.5 yrs	SDR – Alcohol Specific	SDR – Drug Related Causes
Teenage Birth Rate (U20)	519%	SAR – Drug Related Causes	Smoking in Pregnancy
SDR - Drug Related	404%	SAR – Circulatory U75	
Smoking in Pregnancy	367%		

**KEY FINDINGS – SUB-REGIONAL (HSC TRUST & LGD)**

- Male life expectancy either increased or remained similar across the period in all Trusts and Local Government Districts (LGDs), and their most deprived areas, with the exception of the most deprived areas of the South Eastern Trust, Western Trust and Ards & North Down LGD.
- For male life expectancy, the inequality gap between the 20% most deprived areas and the area average widened in the Northern, South Eastern and Western Trusts and the Antrim & Newtownabbey, Ards & North Down and Derry City & Strabane LGDs. The Southern Trust and the Armagh, Banbridge & Craigavon, Mid Ulster and Newry, Mourne & Down LGDs experienced a narrowing of the inequality gap.
- Female life expectancy either increased or remained similar across the period in all Trusts and LGDs, and their most deprived areas. The exception to this was the most deprived areas of the Ards & North Down and Belfast LGDs where it declined.
- For female life expectancy, the inequality gap between the 20% most deprived areas and the area average widened in the Ards & North Down, Armagh City, Banbridge & Craigavon and Belfast LGDs. Conversely, the Antrim & Newtownabbey, Lisburn & Castlereagh, Mid Ulster and Newry, Mourne & Down LGDs experienced a narrowing of their respective inequality gaps.
- Similar to the regional picture, deprivation related inequality was most prominent in indicators relating to alcohol and drugs, self-harm, smoking during pregnancy and teenage births, which were among the five largest inequality gaps for the majority of Trusts and LGDs.
- Drug related mortality was the largest inequality gap in two of the five HSC Trusts and five of the eleven LGDs. In the Western Trust, the rate in its most deprived areas was more than two and a half times (162%) that of the Trust average. While in the Mid & East Antrim LGD the rate in the most deprived areas was more than three times (210%) the LGD average.
- Large inequality gaps for alcohol related admissions also exist in the majority of Trusts and LGDs. In the Western Trust, the rate in its most deprived areas was more than double (128%) that of the Trust average. While in the Derry City & Strabane LGD the rate in the most deprived areas was almost two and half times (139%) the LGD average.
- Deaths due to drug misuse was the largest inequality gap in the Northern (153%) and Western Trusts (177%), with the teenage birth rate the largest inequality gap in the Southern Trust (112%).
- The teenage birth rate was the largest inequality gap in the Armagh City, Banbridge & Craigavon LGD (132%); while alcohol specific mortality showed the largest gap in the Lisburn & Castlereagh LGD (151%).

## INTRODUCTION

This annual publication is one of a series of reports produced as part of the NI Health & Social Care Inequalities Monitoring System (HSCIMS) and presents a comprehensive analysis of health inequality gaps between the most and least deprived areas of NI, and within Health & Social Care (HSC) Trust and Local Government District (LGD) areas across a range of indicators. This report is an update of the Health Inequalities Annual Report 2020. The report is accompanied by downloadable data tables<sup>1</sup> which contain all figures, including urban and rural breakdowns.

The majority of results included in this release are based on figures prior to the coronavirus (COVID-19) pandemic and will not reflect the impact of the pandemic. However, figures related to hospital admissions, emergency care attendances and Primary 1 childhood obesity have been affected to varying degrees. Further details of these will be provided where relevant and included in [Appendix E: Technical Notes & Definitions](#). For information relating to COVID-19 Health Inequalities, please refer to our Coronavirus Health Inequalities Report.<sup>2</sup>





## FORMAT OF THE REPORT

This report is separated into two sections, the first focusing on regional health inequalities and the second presenting sub-regional analysis. The regional section contains separate chapters for each theme/topic area, with each section containing a summary of the key findings, followed by individual indicator analysis. For each indicator two charts are displayed.




## ASSESSMENT OF CHANGE OVER TIME

In addition to the two charts, various symbols are provided that depict changes in the rates in the most deprived and least deprived areas, and in the most-least deprived inequality gap (see below). An indication of the changes observed at the NI level has also been provided for each health outcome. An improvement or decline in the rate is only indicated when the change is statistically significant, or where there is a clear and consistent trend observed over the series. For a notable change in the inequality gap to have occurred, a significant change in at least one of the areas (most/least deprived) has to have been observed, or, where no statistically significant change is apparent then a change in the gap will have deemed to have occurred if there is a clear and consistent trend in both the outcome and the gap over the analysed period. Table 3 overleaf can be used as a reference to aid the reader in understanding how the symbols indicate a change in both the health outcome over time and the resultant inequality gap in this report have been determined.<sup>3</sup>

**Table 1: Indication of change to Indicator Rate**

Changes to indicator rate	
	<p>Most Deprived Areas</p> <p>Least Deprived Areas</p>
	Positive Change
	No Notable Change
	Negative Change

**Table 2: Indication of change in Inequality Gap over time**

Changes in inequality gaps	
	Widening of the gaps
	Narrowing of the gaps
	No notable change in the gap

The sub-regional section presents a condensed summary of findings for each HSC Trust and LGD accompanied by downloadable data tables<sup>2</sup> which contain all figures and an indication of changes to rates and gaps.

<sup>1</sup> <https://www.health-ni.gov.uk/publications/health-inequalities-annual-report-2020>

<sup>2</sup> <https://www.health-ni.gov.uk/articles/coronavirus-related-health-inequalities>

<sup>3</sup> Indicated changes are based on a subjective assessment of the available data.

It should be noted that inequality gaps for indicators can exist in either direction; however health outcomes generally tend to be worse in the most deprived areas than in the least deprived. For the purposes of this report, a positive value for the gap means that the health outcomes experienced in the most deprived areas were worse than in the least deprived.

**Table 3: Understanding changes in the inequality gap**

Change in Health Outcome		Inequality Gap	
Most Deprived Areas	Least Deprived Areas	Symbol	
<b>Gap Widens</b>	Small Positive Change	Large Positive Change	
	Negative Change	Positive Change	
	Negative Change	No Notable Change	
	Large Negative Change	Small Negative Change	
	No Notable Change	Positive Change	
<b>Gap Narrows</b>	Large Positive Change	Small Positive Change	
	Positive Change	Negative Change	
	Positive Change	No Notable Change	
	Small Negative Change	Large Negative Change	
	No Notable Change	Negative Change	
<b>No Notable Change</b>	Positive Change	Positive Change	
	Negative Change	Negative Change	
	No Notable Change	No Notable Change	
	Small Negative Change (Red)/ Positive Change/(Green)	No Notable Change	
	No Notable Change	Small Negative Change (Red)/ Positive Change/(Green)	

Observed differences in the most and least deprived areas, as indicated by the symbol, does not always lead to a change in the gap. Where this has occurred an explanation has been provided where appropriate.



## NOTES FOR USER

- **Regional Inequality Gaps** refer to the difference in health outcomes between the 20% most deprived and 20% least deprived areas of Northern Ireland.
- **Sub-regional Inequality Gaps** refer to the difference between health outcomes for
  - The 20% most deprived areas of an area (LGD/Trust) and the area's average
  - The Trust/LGD and the regional average.
- **Deprivation Measure:** the 20% most and least deprived areas are defined according to the Northern Ireland Multiple Deprivation Measure (NIMDM) 2017.<sup>4</sup>
- **Rounded Figures:** some individual figures have been rounded to either zero or one decimal place independently. As a result, the sum of component items may not therefore always add to the totals shown.
- **Additional Indicators:** figures relating to five additional indicators such as Median Fire Response Times and Median Ambulance Response Times, which form part of the HSCIMS but are not contained in the main body of the report, can be found in Appendix C.
- **Changes to Avoidable Mortality Definitions:** following an Office for National Statistics (ONS) consultation,<sup>5</sup> on a new definition of avoidable mortality as proposed by the Organisation for Economic Co-operation and Development (OECD),<sup>6</sup> a new definition was implemented. Figures based on the old definition have been included in [Appendix C: Additional Indicators](#) for Programme for Government (PfG) monitoring purposes. Full details can be found in [Appendix E: Technical Notes & Definitions](#).
- **Further Analysis:** The appendix section included at the back of the report provides **further analysis** regarding the Social Gradient of Health (Appendix A) and the Population Attributable Risk (PAR) of Deprivation (Appendix B).
- **Urban/Rural Analysis** In addition urban and rural figures for each indicator have been included within the accompanying downloadable tables, and a summary assessment of Rural-NI gaps has been provided in Appendix D.
- **District Electoral Areas (DEAs)** analysis is included within the accompanying downloadable tables. The most recent available health outcomes within each DEA are compared and contrasted with those in the surrounding LGD and notable differences are highlighted.
- **Data limitations** mean that not all health indicators analysed at a regional level can be analysed at Trust, LGD or DEA level. In this report, 45 health indicators have been presented at Trust and 43 at LGD level, with 30 reported at DEA level. A full list can be found in [Table 5](#).
- For **further information** regarding the methodologies, indicator descriptions and sources of data used to produce the analyses throughout this report, please refer to [Appendix E: Technical Notes & Definitions](#).

---

<sup>4</sup> <https://www.nisra.gov.uk/statistics/deprivation/northern-ireland-multiple-deprivation-measure-2017-nimdm2017>

<sup>5</sup> <https://consultations.ons.gov.uk/health-and-life-events/avoidable-mortality-definition/>

<sup>6</sup> <http://www.oecd.org/health/health-systems/Avoidable-mortality-2019-Joint-OECD-Eurostat-List-preventable-treatable-causes-of-death.pdf>

### **Review of suicide statistics in Northern Ireland**

Please note that in light of an on-going review by NISRA and the Coroners' Service into the classification of undetermined deaths, this publication will not use the UK National Statistics definition for suicide. To ensure comparability across years, the sub series relating to self-inflicted injury/intentional self-harm will be used in its place. Full details on this change can be found in [Appendix E: Technical Notes & Definitions](#).

## **Understanding Gaps**

### **Regional Level:**

A positive inequality gap means that the health outcomes in the most deprived areas are worse than in the least deprived areas.

### **Sub-regional Level:**

A positive inequality gap between the Trust or LGD and its most deprived areas means that the health outcomes in the most deprived areas are worse than the Trust or LGD average.

Similarly, a positive inequality gap between the Trust or LGD and NI means that the health outcomes in the Trust or LGD are worse than the NI average.

A negative inequality gap that is widening indicates that the health outcome is experiencing a better change over time within the Trust or LGD than that seen regionally.

Other routine reports in the HSCIMS series include:

**Life Expectancy in Northern Ireland** – presenting the latest official estimates of life expectancy in Northern Ireland and an examination of the causes that contribute to the change in life expectancy over time, as well as the differentials between genders and across Local Government Districts. The latest figures for life expectancy at 65, healthy life expectancy and disability-free life expectancy are also included. <https://www.health-ni.gov.uk/articles/life-expectancy-northern-ireland>

**Making life better: monitoring the wider social determinants of health & wellbeing - key indicators** – monitoring report for the key indicators of the wider social determinants of health & wellbeing, contained in the Making Life Better,<sup>7</sup> the public health strategic framework for NI. <https://www.health-ni.gov.uk/articles/social-determinants-health-statistics>.

**Coronavirus Related Health Inequalities** – presenting an analysis of coronavirus (Covid-19) related health inequalities by assessing differences in positive COVID-19 test cases, COVID-19 admissions and deaths between the most and least deprived areas of Northern Ireland. An assessment of variations across age, sex and urban & rural areas is also included. <https://www.health-ni.gov.uk/articles/coronavirus-related-health-inequalities>

<sup>7</sup> [www.health-ni.gov.uk/topics/public-health-policy-and-advice/making-life-better-whole-system-strategic-framework-public](https://www.health-ni.gov.uk/topics/public-health-policy-and-advice/making-life-better-whole-system-strategic-framework-public)

# Regional Health Inequalities

Regional health inequalities refer to the difference in health outcomes between the 20% most deprived and 20% least deprived areas of Northern Ireland according to the Northern Ireland Multiple Deprivation Measure. This section contains separate chapters for each theme/topic area, with each chapter containing a summary of the key findings, followed by individual indicator analysis.

For each indicator two charts and two symbols are displayed. For ease of understanding, each theme is assigned a separate colour (for example blue is for 'Life Expectancy and General Health'), with a deeper tone representing the 20% most deprived areas and a lighter tone the 20% least deprived.



An example of the indicator analysis, with guidance, can be seen below:

This chart shows trends in rates over time for **Northern Ireland**, the **20% most deprived areas** and **20% least deprived areas**.

This symbol for assessment of change over time is explained on pages 7 and 8 of this report. Also shown is an indication of change at the Northern Ireland (NI) level.

## Male Life Expectancy at Birth

NI 



This chart shows the trend for the most-least deprived inequality gap over the same period. The gap between most-least deprived may be displayed as a discrete value, e.g. years for life expectancies or as a percentage difference.

SUMMARY OF CHANGES IN REGIONAL INEQUALITY GAPS OVER THE LAST 5 YEARS<sup>8</sup>

< > **Most-Least Deprived Inequality Gaps that Widened over the Analysed Period**

**9** indicators had inequality gaps that **widened** over the period analysed

	Change in Health Outcome		
	Northern Ireland	Most Deprived Areas	Least Deprived Areas
Potential Years of Life Lost	Positive Change	Positive Change	Positive Change
Standardised Death Rate - Preventable	No Notable Change	No Notable Change	Positive Change
Standardised Prescription Rate - Statin	Positive Change	No Notable Change	Positive Change
Standardised Prescription Rate - Mood & Anxiety	Negative Change	Negative Change	Negative Change
Standardised Death Rate - Drug Related Causes	Negative Change	Negative Change	Negative Change
Standardised Death Rate - Drug Misuse	Negative Change	Negative Change	Negative Change
Infant Mortality	No Notable Change	No Notable Change	Positive Change
Smoking During Pregnancy	Positive Change	No Notable Change	Positive Change
Low Birth Weight	No Notable Change	No Notable Change	Positive Change

**Key:** Negative Change No Notable Change Positive Change

> < **Most-Least Deprived Inequality Gaps that Narrowed over the Analysed Period**

**12** indicators had inequality gaps that **narrowed** over the period analysed

	Change in Health Outcome		
	Northern Ireland	Most Deprived Areas	Least Deprived Areas
Standardised Death Rate - Treatable	Positive Change	Positive Change	No Notable Change
Standardised Admissions Rate - Circulatory	Positive Change	Positive Change	Positive Change
Standardised Admission Rate - Circulatory U75	Positive Change	Positive Change	Positive Change
Standardised Incidence Rate - Cancer	Negative Change	Negative Change	Negative Change
Standardised Admission Rate - All Admissions	Positive Change	Positive Change	Positive Change
Standardised Admission Rate - Emergency Admissions	Positive Change	Positive Change	Positive Change
Standardised Admission Rate - Self-Harm	Positive Change	Positive Change	Positive Change
Crude Death Rate - Intentional Self-Harm	Positive Change	Positive Change	No Notable Change
Standardised Admission Rate - Alcohol Related Causes	Positive Change	Positive Change	Negative Change
Standardised Death Rate - Alcohol Specific Causes	Negative Change	Negative Change	Negative Change
Standardised Admission Rate - Drug Related Causes	Positive Change	Positive Change	Positive Change
Breastfeeding on Discharge	Positive Change	Positive Change	Positive Change

**Key:** Negative Change No Notable Change Positive Change

<sup>8</sup> There are some indicators, Small for Gestational Age and Year 8 Obesity, for which a regional assessment of change has not been carried out. Further details are provided alongside each indicator where relevant.

## Most-Least Deprived Inequality Gaps that Showed No Notable Change over the Analysed Period

**27** indicators had inequality gaps that **showed no notable change** over the period analysed

	Change in Health Outcome		
	Northern Ireland	Most Deprived Areas	Least Deprived Areas
Male Life Expectancy at Birth	Positive Change	Positive Change	Positive Change
Female Life Expectancy at Birth	Positive Change	No Notable Change	No Notable Change
Male Life Expectancy at Age 65	Positive Change	No Notable Change	Positive Change
Female Life Expectancy at Age 65	Positive Change	No Notable Change	No Notable Change
Male Healthy Life Expectancy	No Notable Change	No Notable Change	No Notable Change
Female Healthy Life Expectancy	No Notable Change	No Notable Change	No Notable Change
Male Disability Free Life Expectancy	No Notable Change	No Notable Change	No Notable Change
Female Disability Free Life Expectancy	No Notable Change	No Notable Change	No Notable Change
Standardised Death Rate – Avoidable	Positive Change	Positive Change	Positive Change
Standardised Death Rate – Avoidable: Children & Young People	No Notable Change	No Notable Change	No Notable Change
Standardised Death Rate – Circulatory U75	Positive Change	Positive Change	Positive Change
Standardised Death Rate – Respiratory U75	No Notable Change	Negative Change	No Notable Change
Standardised Death Rate - Cancer U75	Positive Change	Positive Change	Positive Change
Standardised Death Rate - All Cause U75	Positive Change	Positive Change	Positive Change
Standardised Prescription Rate - Anti hypertensive	Positive Change	Positive Change	Positive Change
Standardised Admission Rate - Respiratory	Positive Change	Positive Change	Positive Change
Standardised Admission Rate - Respiratory U75	Positive Change	Positive Change	Positive Change
Standardised Attendance Rate - Emergency Care	Negative Change	Negative Change	Negative Change
Standardised Admission Rate - Elective Inpatient Admissions	Positive Change	Positive Change	Positive Change
Standardised Admission Rate – Day Case	Positive Change	Positive Change	Positive Change
Standardised Death Rate - Smoking Related Causes	Positive Change	Positive Change	Positive Change
Standardised Incidence Rate - Lung Cancer	Negative Change	No Notable Change	No Notable Change
Standardised Death Rate - Lung Cancer	No Notable Change	Positive Change	No Notable Change
Teenage Birth Rate U20	Positive Change	No Notable Change	No Notable Change
Healthy Birth Weight	No Notable Change	No Notable Change	No Notable Change
Primary 1 BMI: Obese	No Notable Change	No Notable Change	No Notable Change
Primary 1 BMI: Overweight or Obese	No Notable Change	No Notable Change	No Notable Change

**Key:**

Negative Change
No Notable Change
Positive Change

## Life Expectancy & General Health



In 2017-19 the life expectancy gender gap between males and females in Northern Ireland was 3.8 years. As with all life expectancy and general health indicators analysed there was no change in the deprivation gap for male life expectancy at birth, although life expectancy improved across all areas. There was a positive change in female life expectancy at birth overall, although no change in the most and least deprived areas. Life expectancy at 65 increased for both genders, however there was no change for both healthy life expectancy and disability free life expectancy for males or females.

**Key:** Northern Ireland: NI




**MD** Most Deprived Areas  
**LD** Least Deprived Areas

Positive Change }< Narrowing of the Gap  
 No Notable Change — No Notable Change in the Gap  
 Negative Change }> Widening of the Gap



### Male Life Expectancy at Birth NI



### Female Life Expectancy at Birth NI





### Male Life Expectancy at Age 65 NI





Female Life Expectancy at Age 65

NI 





Male Healthy Life Expectancy

NI 





Female Healthy Life Expectancy

NI 



Male Disability Free Life Expectancy<sup>9</sup>



NI 



<sup>9</sup> A change in the wording of the Health Survey NI question in 2012/13 from which the DFLE indicators are derived may partly explain any observed decreases in DFLE estimates. Further information is included under 'Indicator Definitions' within the Appendices.

Female Disability Free Life Expectancy<sup>9</sup>


NI 







## Premature Mortality<sup>10</sup>


Rates of premature mortality generally decreased over the period in NI and its most and least deprived areas. However, large inequality gaps continue to persist with the most deprived areas continuing to experience higher mortality rates than the least deprived areas. The inequality gaps remained similar across all indicators with the exception of deaths from preventable causes where the gap widened due to decreased mortality in the least deprived areas and treatable mortality where the gap narrowed. For respiratory mortality among under 75s, the rate in the most deprived areas was three and a half times that seen in the least deprived.





### Potential Years of Life Lost



### Standardised Death Rate - Treatable<sup>10</sup>





### Standardised Death Rate – Preventable<sup>10</sup>



<sup>10</sup> Based on the new OECD definition of avoidable mortality. The previous definition used for the Programme for Government indicator for preventable mortality will continue to be calculated using the old definition and is published in [Appendix C](#). A full explanation, including indicator definitions, can be found in [Appendix E](#).



Standardised Death Rate – Avoidable<sup>11</sup>

NI 




Standardised Death Rate – Avoidable: Children & Young People<sup>12</sup>

NI 






Standardised Death Rate – Circulatory U75

NI 



Standardised Death Rate – Respiratory U75

NI 





<sup>11</sup> Based on the new OECD definition of avoidable mortality. The previous definition used for the Programme for Government indicator for preventable mortality will continue to be calculated using the old definition and is published in [Appendix C](#). A full explanation, including indicator definitions, can be found in [Appendix E](#).

<sup>12</sup> This indicator should be treated as an experimental statistic. Given the relatively small numbers and large variability in specific causes of childhood deaths, interpretation should be made with caution.



Standardised Death Rate – Cancer U75

NI




Standardised Death Rate – All Cause U75

NI








## Major Diseases<sup>13</sup>




The most-least deprived inequality gap remained similar or narrowed for all indicators with the exception of the prescription rate for statins. There were improvements in all indicators at a regional level, with the exception of cancer incidence which increased over the period across all areas. The largest inequality gap was observed for admissions due to respiratory diseases, with the admission rate in the most deprived areas remaining around double that of the least deprived areas, for all ages and for those aged under 75 years.

**Key:** Northern Ireland: NI 



**MD** Most Deprived Areas  
**LD** Least Deprived Areas

 Positive Change      Narrowing of the Gap



 No Notable Change     No Notable Change in the Gap

 Negative Change      Widening of the Gap



### Standardised Admission Rate – Circulatory NI



### Standardised Admission Rate – Circulatory U75 NI






### Standardised Prescription Rate – Antihypertensive NI



<sup>13</sup> Mental health related conditions, alcohol and drug related conditions, are considered in separate chapters.



Standardised Prescription Rate – Statin

NI 





Standardised Admission Rate – Respiratory

NI 






Standardised Admission Rate – Respiratory U75

NI 



Standardised Incidence Rate – Cancer

NI 



## Hospital Activity<sup>14</sup>



Inequality gaps for all indicators remained constant over the period with the exception of all admissions and emergency admissions, which narrowed. Emergency care attendances was the only indicator to see a negative change across NI and its most and least deprived areas. All other hospital activity indicators improved across NI and in its most and least deprived areas. Even though the emergency admissions inequality gap narrowed, it remained the largest of the four indicators analysed at three-fifths higher in the most deprived areas.

**Key:** Northern Ireland: NI



**MD** Most Deprived Areas  
**LD** Least Deprived Areas

Positive Change      Narrowing of the Gap  
 No Notable Change      No Notable Change in the Gap  
 Negative Change      Widening of the Gap



### Standardised Admission Rate – All Admissions NI



### Standardised Admission Rate – Emergency Admissions NI





### Standardised Attendance Rate – Emergency Care NI



<sup>14</sup> Please note that hospital activity figures for 2019/20 may have been affected by changes in activity levels as a result of the COVID-19 pandemic. Further detail can be found in [Appendix E: Technical Notes & Definitions](#).



Standardised Admission Rate – Elective Inpatient Admissions<sup>15</sup>

NI



Standardised Admission Rate – Day Case Admissions

NI



<sup>15</sup> Please note that some of the observed decreases in the admission rate across all areas may be explained by a service reconfiguration. For further information, see indicator definitions in [Appendix E: Technical Notes & Definitions](#).

## Mental Health



Large inequality gaps continue to exist for mental health indicators, with the latest position showing that the death rate from intentional self-harm in the most deprived areas was double that in the least deprived areas. There was positive change regionally and in the most and least deprived areas for admissions due to self-harm, with the inequality gap also narrowing.<sup>16</sup> Prescription rates for mood and anxiety disorders increased in NI and its most and least deprived areas, with the gap widening.

**Key:** Northern Ireland: NI



**MD** Most Deprived Areas  
**LD** Least Deprived Areas

Positive Change      Narrowing of the Gap  
 No Notable Change      No Notable Change in the Gap  
 Negative Change      Widening of the Gap



### Standardised Admission Rate – Self-Harm<sup>16</sup>



### Crude Death Rate – Intentional Self-Harm<sup>17</sup>



### Standardised Prescription Rate – Mood & Anxiety




<sup>16</sup> Please note that some of the observed decreases in the admission rate across all areas may be explained by a service reconfiguration. For further information, see indicator definitions in [Appendix E: Technical Notes & Definitions](#).

<sup>17</sup> This indicator has replaced Crude Death Rate - Suicide. Full details on this change can be found in [Appendix E](#).









## Alcohol, Smoking & Drugs

Alcohol, smoking and drug related indicators continued to have some of the largest health inequalities in NI. The inequality gap for drug related and drug misuse deaths widened, while the gaps narrowed or remained similar for other indicators. Alcohol specific mortality and alcohol related admissions in the most deprived areas were around four times that in the least deprived areas. Although lung cancer incidence rose in NI overall, the inequality gap remained unchanged. While the drug related admission rate fell across all areas, the opposite was true for drug related death rates and drug misuse.<sup>18</sup>



**Key:** Northern Ireland: NI 

**MD** Most Deprived Areas  
**LD** Least Deprived Areas

 Positive Change       Narrowing of the Gap  
 No Notable Change       No Notable Change in the Gap  
 Negative Change       Widening of the Gap




### Standardised Admission Rate – Alcohol Related Causes

NI 





### Standardised Death Rate – Alcohol Specific

NI 



### Standardised Death Rate – Smoking Related Causes



NI 



<sup>18</sup> Please note that some observations may be due to changes in drug misuse behaviours among the population. For more information see [Appendix E: Technical Notes & Definitions](#).



Standardised Incidence Rate – Lung Cancer

NI 





Standardised Death Rate – Lung Cancer

NI 





Standardised Admission Rate – Drug Related Causes

NI 





Standardised Death Rate – Drug Related Causes

NI 






Standardised Death Rate – Drug Misuse









## Pregnancy & Early Years

Of all the indicators related to pregnancy and early years, only breastfeeding on discharge showed a narrowing of the inequality gap. The low birth weight gap widened, due to improvements in the least deprived areas. This was also true for the gap between the most and least deprived areas for smoking during pregnancy. The inequality gaps for the under 20 teenage birth rate and the proportion of mothers smoking during pregnancy still remain very large. The teenage birth rate in the most deprived areas was more than six times that in the least deprived areas.



**Key:** Northern Ireland: NI 

 **MD** Most Deprived Areas  
 **LD** Least Deprived Areas

 Positive Change  Narrowing of the Gap  
 No Notable Change  No Notable Change in the Gap  
 Negative Change  Widening of the Gap




### Infant Mortality Rate<sup>19</sup>

NI 





### Smoking During Pregnancy

NI 



### Teenage Birth Rate U20



NI 



<sup>19</sup> As the underlying rates are relatively low (typically below 6 deaths per 1,000 live births), small annual changes can have a large impact on the resulting inequality gap.



Breastfeeding on Discharge

NI 





Low Birth Weight

NI 




Healthy Birth Weight

NI 



Small for Gestational Age<sup>20</sup>



<sup>20</sup> No assessment of the inequality gap trend will be made until 5 years of data are available.


## Childhood Obesity

Over the period analysed there was no notable change in the proportion of Primary 1 children reported as overweight or obese.<sup>21</sup> However, rates of obesity are continually higher in the most deprived areas.

Please note that since 2018/19, Year 8 figures for Northern Ireland do not include measurements from the Western HSC Trust, therefore no assessment of change relating to NI values will be provided.<sup>22</sup> In addition, Primary 1 figures combine 3 years of data, as the number of measurements recorded in 2019/20 was impacted by school closures due to the COVID-19 pandemic.









**Key:**

Northern Ireland: NI 





MD Most Deprived Areas

LD Least Deprived Areas

-  Positive Change
-  No Notable Change
-  Negative Change
-   Narrowing of the Gap
-  No Notable Change in the Gap
-   Widening of the Gap



### Primary 1 BMI: Obese

NI 





### Primary 1 BMI: Overweight or Obese

NI 





### Year 8 BMI: Obese<sup>22</sup>



<sup>21</sup> As the underlying figures are somewhat low, small annual changes can have a large impact on the observed inequality gap.  
<sup>22</sup> No assessment of Year 8 Obesity will be made as data for the Western Trust since 2018/19 is unavailable. A full explanation of any data issues can be found in [Appendix E: Technical Notes & Definitions](#).

Year 8 BMI: Overweight or Obese<sup>23</sup>



<sup>23</sup> No assessment of Year 8 Obesity will be made as data for the Western Trust since 2018/19 is unavailable. A full explanation of any data issues can be found in [Appendix E: Technical Notes & Definitions](#).

# Sub-regional Health Inequalities

There are two aspects to the analysis of sub-regional health inequalities in this section. The analysis for each sub-regional area will either relate to the difference in health outcomes between the **Trust or LGD and the regional (NI) average** or to the **20% most deprived areas within an area and the area's average**.

Each chapter is a summary of findings only. For a full assessment of the HSC Trust and LGD figures, including a range of indicators that are also available at District Electoral Area (DEA), see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

An example of a sub-regional analysis, with guidance, can be seen below:

Each chapter contains a comparison of the area's health outcomes against the regional average including a summary of the most notable indicators that were better or worse.

There is also a comparison of the Inequality Gap between the area average and its most deprived areas in respect of:

**Life Expectancy**


Analysis of the male and female life expectancy gap within the area and if this gap has changed.

**Largest Inequality Gaps**

The five largest inequality gaps in the area.

**Changes in Inequality Gaps**

A summary of the most notable indicators that have either narrowed or widened across the analysed period.






## SUMMARY OF CHANGES IN SUB-REGIONAL INEQUALITY GAPS OVER THE LAST 5 YEARS

## Changes in Deprivation Related Inequality Gaps

Over the period analysed, there were more inequality gaps that widened than narrowed in the Northern, South Eastern and Southern HSC Trusts, with an equal amount in the Belfast and Western HSC Trusts. Changes in the inequality gaps varied across LGDs with 16 indicators widening in Ards & North Down and 17 indicators narrowing in Newry, Mourne and Down.

For each area analysed, the chart below shows the number of indicators that widened, narrowed or did not show a notable change across the period.



\* For the purposes of this graphic, gaps which reversed direction, but remained similar in magnitude have been included in the "No Change" category.


**COMPARISON OF SUB-REGIONAL HEALTH OUTCOMES AGAINST THE REGIONAL AVERAGE**

The following areas had a majority of health outcomes that were better than the NI average:

- South Eastern HSC Trust
- Ards & North Down
- Lisburn & Castlereagh
- Mid Ulster

The following areas had a majority of health outcomes that were worse than the NI average:

- Belfast HSC Trust
- Belfast LGD
- Derry City & Strabane



## Largest Deprivation Inequality Gaps in Each Area


The table below indicates the five largest deprivation inequality gaps in each Health & Social Care Trust (HSCT) and Local Government District (LGD).

<b>Belfast HSCT</b>	SDR Drug Related (116%)	SDR Drug Misuse (111%)	SAR Alcohol Related (107%)	SDR Respiratory U75 (104%)	SAR Drug Related (99%)
<b>Northern HSCT</b>	SDR Drug Misuse (153%)	SDR Drug Related (151%)	SDR Alcohol Specific (108%)	SAR Drug Related (101%)	Teenage Birth Rate U20 (98%)
<b>South Eastern HSCT</b>	SDR Drug Related (136%)	SDR Drug Misuse (135%)	Smoking During Pregnancy (101%)	Teenage Birth Rate (U20) (98%)	SDR Alcohol Specific (97%)
<b>Southern HSCT</b>	Teenage Birth Rate (U20) (112%)	SAR Alcohol Related (106%)	SAR Drug Related (95%)	SAR Self-Harm (92%)	SDR Respiratory U75 (87%)
<b>Western HSCT</b>	SDR Drug Misuse (177%)	SDR Drug Related (162%)	SAR Alcohol Related (128%)	SDR Alcohol Specific (113%)	SAR Self-Harm (101%)
<b>Antrim &amp; Newtownabbey LGD</b>	SDR Drug Related (147%)	SDR Alcohol Specific (141%)	Teenage Birth Rate (U20) (141%)	SAR Drug Related (97%)	SAR Self-Harm (96%)
<b>Ards &amp; North Down LGD</b>	SDR Drug Related (114%)	SDR Alcohol Specific (102%)	Smoking During Pregnancy (100%)	Teenage Birth Rate (U20) (96%)	CDR Intentional Self-Harm (86%)
<b>Armagh City, Banbridge &amp; Craigavon LGD</b>	Teenage Birth Rate (U20) (132%)	SAR Alcohol Related (116%)	SAR Drug Related (114%)	SDR Alcohol Specific (101%)	SAR Self-Harm (90%)
<b>Belfast LGD</b>	SAR Alcohol Related (96%)	SAR Drug Related (89%)	SAR Self-Harm (88%)	SDR Drug Related (86%)	Teenage Birth Rate (U20) (78%)
<b>Causeway Coast &amp; Glens LGD</b>	SDR Drug Related (170%)	SDR Alcohol Specific (130%)	SAR Alcohol Related (112%)	SAR Self-Harm (84%)	CDR Intentional Self-Harm (76%)
<b>Derry City &amp; Strabane LGD</b>	SAR Alcohol Related (139%)	SDR Alcohol Specific (131%)	SAR Self-Harm (112%)	Teenage Birth Rate (U20) (104%)	SDR Drug Related (101%)
<b>Fermanagh &amp; Omagh LGD</b>	SAR Drug Related (78%)	Teenage Birth Rate (U20) (74%)	SDR Alcohol Specific (61%)	SAR Self-Harm (59%)	SDR Drug Related (59%)
<b>Lisburn &amp; Castlereagh LGD</b>	SDR Alcohol Specific (151%)	Smoking During Pregnancy (136%)	Teenage Birth Rate (U20) (120%)	SDR Lung Cancer (93%)	SAR Alcohol Related (93%)
<b>Mid &amp; East Antrim LGD</b>	SDR Drug Related (210%)	SAR Drug Related (139%)	SAR Self-Harm (136%)	SAR Alcohol Related (132%)	Teenage Birth Rate (U20) (131%)
<b>Mid Ulster LGD</b>	Teenage Birth Rate (U20) (72%)	SDR Alcohol Specific (68%)	SAR Self-Harm (59%)	SAR Alcohol Related (58%)	SAR Drug Related (56%)
<b>Newry, Mourne &amp; Down LGD</b>	SDR Drug Related (90%)	Teenage Birth Rate (U20) (65%)	SAR Self-Harm (61%)	SAR Alcohol Related (60%)	SAR Drug Related (52%)

# Belfast

## Health & Social Care Trust (HSCT)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Belfast Trust and its 20% most deprived areas:

#### Life Expectancy

- Male life expectancy in the Trust’s most deprived areas was 70.9 years, **5.7 years** less than the HSCT average (76.6 years). No Change in Male Inequality Gap
- Female life expectancy in the Trust’s most deprived areas was 77.1 years, **4.2 years** less than the HSCT average (81.2 years). No Change in Female Inequality Gap

#### Largest Inequality Gaps

SDR Drug Related (116%)	SDR Drug Misuse (111%)	SAR Alcohol Related (107%)	SDR Respiratory U75 (104%)	SAR Drug Related (99%)
-------------------------	------------------------	----------------------------	----------------------------	------------------------


#### Changes in Inequality Gaps

Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Elective Inpatient Admissions Crude Intentional Self-Harm Rate Day Case Admissions	Primary 1 Overweight or Obesity Drug Misuse Smoking During Pregnancy

This is a summary of findings only. For a full assessment and all figures see downloadable tables at: <https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Northern Health & Social Care Trust (HSCT)

## Comparison with NI



## Comparison with Most Deprived Areas

Inequality gaps between the Northern Trust and its 20% most deprived areas:


### Life Expectancy

- Male life expectancy in the Trust’s most deprived areas was 75.8 years, **3.7 years** less than the HSCT average (79.5 years). **Male Inequality Gap Widened**
- Female life expectancy in the Trust’s most deprived areas was 80.4 years, **2.5 years** less than the HSCT average (82.9 years). **No Change in Female Inequality Gap**

### Largest Inequality Gaps

SDR Drug Misuse (153%)	SDR Drug Related (151%)	SDR Alcohol Specific (108%)	SAR Drug Related (101%)	Teenage Birth Rate U20 (98%)
------------------------	-------------------------	-----------------------------	-------------------------	------------------------------

### Changes in Inequality Gaps




This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# South Eastern Health & Social Care Trust (HSCT)

## Comparison with NI



## Comparison with Most Deprived Areas

Inequality gaps between the South Eastern Trust and its 20% most deprived areas:

### Life Expectancy

- Male Inequality Gap Widened
- No Change in Female Inequality Gap

### Largest Inequality Gaps

SDR Drug Related (136%)	SDR Drug Misuse (135%)	Smoking During Pregnancy (101%)	Teenage Birth Rate (U20) (98%)	SDR Alcohol Specific (97%)
-------------------------	------------------------	---------------------------------	--------------------------------	----------------------------

### Changes in Inequality Gaps


<b>Most Notable Narrowed</b> Deprivation Inequality Gaps	<b>Most Notable Widened</b> Deprivation Inequality Gaps
Crude Intentional Self-Harm Rate Alcohol Specific Mortality Self-Harm Admissions	Cancer/Lung Cancer Incidence Drug Related/Misuse Mortality Statin Prescription Rate

**This is a summary of findings only. For a full assessment and all figures see downloadable tables at:**  
<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Southern

## Health & Social Care Trust (HSCT)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Southern Trust and its 20% most deprived areas:

#### Life Expectancy

- Male life expectancy in the Trust's most deprived areas was 77.1 years, **2.2 years** less than the HSCT average (79.3 years). Male Inequality Gap Narrowed
- Female life expectancy in the Trust's most deprived areas was 81.6 years, **1.5 years** less than the HSCT average (83.1 years). No Change in Female Inequality Gap

#### Largest Inequality Gaps

Teenage Birth Rate (U20) (112%)	SAR Alcohol Related (106%)	SAR Drug Related (95%)	SAR Self-Harm (92%)	SDR Respiratory U75 (87%)
---------------------------------	----------------------------	------------------------	---------------------	---------------------------

#### Changes in Inequality Gaps

Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Male Life Expectancy at Birth Breastfeeding on Discharge Alcohol Specific Mortality	Low Birth Weight Primary 1 Obesity Teenage Birth Rate (U20)


This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Western

## Health & Social Care Trust (HSCT)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Western Trust and its 20% most deprived areas:


#### Life Expectancy

- Male life expectancy in the Trust’s most deprived areas was 74.1 years, **4.4 years** less than the HSCT average (78.5 years). **Male Inequality Gap Widened**
- Female life expectancy in the Trust’s most deprived areas was 79.4 years, **2.8 years** less than the HSCT average (82.2 years). **No Change in Female Inequality Gap**

#### Largest Inequality Gaps

SDR Drug Misuse (177%)	SDR Drug Related (162%)	SAR Alcohol Related (128%)	SDR Alcohol Specific (113%)	SAR Self-Harm (101%)
------------------------	-------------------------	----------------------------	-----------------------------	----------------------

#### Changes in Inequality Gaps



**This is a summary of findings only. For a full assessment and all figures see downloadable tables at:**


<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>



# Antrim & Newtownabbey

## Local Government District (LGD)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Antrim & Newtownabbey LGD and its 20% most deprived areas:

#### Life Expectancy

- Male life expectancy in the LGD's most deprived areas was 75.2 years, **4.3 years** less than the LGD average (79.5 years). **Male Inequality Gap Widened**
- Female life expectancy in the LGD's most deprived areas was 79.2 years, **3.4 years** less than the LGD average (82.7 years). **Female Inequality Gap Narrowed**

#### Largest Inequality Gaps

SDR Drug Related (147%)	SDR Alcohol Specific (141%)	Teenage Birth Rate (U20) (141%)	SAR Drug Related (97%)	SAR Self-Harm (96%)
----------------------------	--------------------------------	------------------------------------	---------------------------	------------------------

#### Changes in Inequality Gaps

Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Elective Inpatient Admissions U75 Circulatory Admissions Respiratory Admissions	Crude Intentional Self-Harm Rate Male Life Expectancy at 65 Alcohol Specific Mortality


This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Ards & North Down

## Local Government District (LGD)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Ards & North Down LGD and its 20% most deprived areas:

#### Life Expectancy

- Male life expectancy in the LGD's most deprived areas was 75.9 years, **3.7 years** less than the LGD average (79.5 years). Male Inequality Gap Widened
- Female life expectancy in the LGD's most deprived areas was 79.8 years, **3.1 years** less than the LGD average (82.9 years). Female Inequality Gap Widened

#### Largest Inequality Gaps

SDR Drug Related (114%)	SDR Alcohol Specific (102%)	Smoking During Pregnancy (100%)	Teenage Birth Rate (U20) (96%)	CDR Intentional Self-Harm (86%)
-------------------------	-----------------------------	---------------------------------	--------------------------------	---------------------------------

#### Changes in Inequality Gaps

Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Low Birth Weight Primary 1 Obesity/Overweight Emergency Attendances	Crude Intentional Self-Harm Rate Elective Inpatient Admissions Cancer Incidence


This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Armagh City, Banbridge & Craigavon

## Local Government District (LGD)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Armagh City, Banbridge & Craigavon LGD and its 20% most deprived areas:

#### Life Expectancy

- Male life expectancy in the LGD's most deprived areas was 76.3 years, **3.2 years** less than the LGD average (79.5 years). Male Inequality Gap Narrowed
- Female life expectancy in the LGD's most deprived areas was 81.5 years, **1.6 years** less than the LGD average (83.1 years). Female Inequality Gap Widened

#### Largest Inequality Gaps

Teenage Birth Rate (U20) (132%)	SAR Alcohol Related (116%)	SAR Drug Related (114%)	SDR Alcohol Specific (101%)	SAR Self-Harm (90%)
---------------------------------	----------------------------	-------------------------	-----------------------------	---------------------

#### Changes in Inequality Gaps


Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Treatable Mortality U75 Circulatory Mortality Lung Cancer Mortality	Low Birth Weight Male Life Expectancy at 65 Teenage Birth Rate (U20)

This is a summary of findings only. For a full assessment and all figures see downloadable tables at: <https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Belfast

## Local Government District (LGD)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Belfast LGD and its 20% most deprived areas:


#### Life Expectancy

- Male life expectancy in the LGD’s most deprived areas was 71.3 years, **4.7 years** less than the LGD average (76.1 years). **No Change in Male Inequality Gap**
- Female life expectancy in the LGD’s most deprived areas was 77.0 years, **4.0 years** less than the LGD average (81.0 years). **Female Inequality Gap Widened**

#### Largest Inequality Gaps

SAR Alcohol Related (96%)	SAR Drug Related (89%)	SAR Self-Harm (88%)	SDR Drug Related (86%)	Teenage Birth Rate (U20) (78%)
---------------------------	------------------------	---------------------	------------------------	--------------------------------

#### Changes in Inequality Gaps




This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Causeway Coast & Glens

## Local Government District (LGD)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Causeway Coast & Glens LGD and its 20% most deprived areas:

#### Life Expectancy

- Male life expectancy in the LGD's most deprived areas was 75.9 years, **3.5 years** less than the LGD average (79.5 years). No Change in Male Inequality Gap
- Female life expectancy in the LGD's most deprived areas was 81.3 years, **1.6 years** less than the LGD average (82.9 years). No Change in Female Inequality Gap

#### Largest Inequality Gaps

SDR Drug Related (170%)	SDR Alcohol Specific (130%)	SAR Alcohol Related (112%)	SAR Self-Harm (84%)	CDR Intentional Self-Harm (76%)
----------------------------	--------------------------------	-------------------------------	------------------------	------------------------------------


#### Changes in Inequality Gaps

Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Elective Inpatient Admissions Treatable Mortality U75 Circulatory Mortality	Primary 1 Obesity/Overweight Circulatory Admissions Alcohol Specific Mortality

This is a summary of findings only. For a full assessment and all figures see downloadable tables at: <https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Derry City & Strabane Local Government District (LGD)

## Comparison with NI



## Comparison with Most Deprived Areas

Inequality gaps between the Derry City & Strabane LGD and its 20% most deprived areas:


### Life Expectancy

- Male life expectancy in the LGD’s most deprived areas was 72.5 years, **5.3 years** less than the LGD average (77.8 years). **Male Inequality Gap Widened**
- Female life expectancy in the LGD’s most deprived areas was 78.5 years, **2.9 years** less than the LGD average (81.4 years). **No Change in Female Inequality Gap**

### Largest Inequality Gaps

SAR Alcohol Related (139%)	SDR Alcohol Specific (131%)	SAR Self-Harm (112%)	Teenage Birth Rate (U20) (104%)	SDR Drug Related (101%)
----------------------------	-----------------------------	----------------------	---------------------------------	-------------------------

### Changes in Inequality Gaps




This is a summary of findings only. For a full assessment and all figures see downloadable tables at: <https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Fermanagh & Omagh

## Local Government District (LGD)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Fermanagh & Omagh LGD and its 20% most deprived areas:

#### Life Expectancy

- Male life expectancy in the LGD's most deprived areas was 77.9 years, **1.4 years** less than the LGD average (79.3 years). No Change in Male Inequality Gap
- Female life expectancy in the LGD's most deprived areas was 81.1 years, **2.0 years** less than the LGD average (83.2 years). No Change in Female Inequality Gap

#### Largest Inequality Gaps

SAR Drug Related (78%)	Teenage Birth Rate (U20) (74%)	SDR Alcohol Specific (61%)	SAR Self-Harm (59%)	SDR Drug Related (59%)
------------------------	--------------------------------	----------------------------	---------------------	------------------------


#### Changes in Inequality Gaps

Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Primary 1 Obesity/Overweight Alcohol Related Admissions Emergency Admissions	Lung Cancer Incidence Teenage Birth Rate (U20) Cancer Incidence

This is a summary of findings only. For a full assessment and all figures see downloadable tables at: <https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Lisburn & Castlereagh Local Government District (LGD)

## Comparison with NI



## Comparison with Most Deprived Areas

Inequality gaps between the Lisburn & Castlereagh LGD and its 20% most deprived areas:


### Life Expectancy

- Male life expectancy in the LGD’s most deprived areas was 76.1 years, **4.1 years** less than the LGD average (80.1 years). **No Change in Male Inequality Gap**
- Female life expectancy in the LGD’s most deprived areas was 81.8 years, **1.7 years** less than the LGD average (83.5 years). **Female Inequality Gap Narrowed**

### Largest Inequality Gaps

SDR Alcohol Specific (151%)	Smoking During Pregnancy (136%)	Teenage Birth Rate (U20) (120%)	SDR Lung Cancer (93%)	SAR Alcohol Related (93%)
-----------------------------	---------------------------------	---------------------------------	-----------------------	---------------------------

### Changes in Inequality Gaps




This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>



# Mid and East Antrim Local Government District (LGD)

## Comparison with NI



## Comparison with Most Deprived Areas

Inequality gaps between the Mid and East Antrim LGD and its 20% most deprived areas:

### Life Expectancy

- Male life expectancy in the LGD’s most deprived areas was 74.7 years, **4.4 years** less than the LGD average (79.1 years). No Change in Male Inequality Gap
- Female life expectancy in the LGD’s most deprived areas was 79.1 years, **3.6 years** less than the LGD average (82.7 years). No Change in Female Inequality Gap

### Largest Inequality Gaps

SDR Drug Related (210%)	SAR Drug Related (139%)	SAR Self-Harm (136%)	SAR Alcohol Related (132%)	Teenage Birth Rate (U20) (131%)
-------------------------	-------------------------	----------------------	----------------------------	---------------------------------

### Changes in Inequality Gaps


Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Crude Intentional Self-Harm Rate Antihypertensive Prescriptions U75 Respiratory Admissions	Low Birth Weight Primary 1 Obesity Teenage Birth Rate (U20)

This is a summary of findings only. For a full assessment and all figures see downloadable tables at: <https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Mid Ulster

## Local Government District (LGD)

### Comparison with NI



### Comparison with Most Deprived Areas

Inequality gaps between the Mid Ulster LGD and its 20% most deprived areas:

#### Life Expectancy

- Male life expectancy in the LGD’s most deprived areas was 77.9 years, 1.8 years less than the LGD average (79.7 years).
Male Inequality Gap Narrowed
- Female life expectancy in the LGD’s most deprived areas was 83.1 years, 0.1 years less than the LGD average (83.2 years).
Female Inequality Gap Narrowed

#### Largest Inequality Gaps

Teenage Birth Rate (U20) (72%)	SDR Alcohol Specific (68%)	SAR Self-Harm (59%)	SAR Alcohol Related (58%)	SAR Drug Related (56%)
--------------------------------	----------------------------	---------------------	---------------------------	------------------------

#### Changes in Inequality Gaps


Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Female Life Expectancy at Birth Crude Intentional Self-Harm Rate Smoking During Pregnancy	Male Life Expectancy at 65 Alcohol Specific Mortality Elective Inpatient Admissions

This is a summary of findings only. For a full assessment and all figures see downloadable tables at:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

# Newry, Mourne and Down Local Government District (LGD)

## Comparison with NI



## Comparison with Most Deprived Areas

Inequality gaps between the Newry, Mourne & Down LGD and its 20% most deprived areas:

### Life Expectancy

- Male life expectancy in the LGD’s most deprived areas was 78.1 years, **1.1 years** less than the LGD average (79.2 years).
 

 Male Inequality Gap  
Narrowed
- Female life expectancy in the LGD’s most deprived areas was 82.4 years, **0.9 years** less than the LGD average (83.3 years).
 

 Female Inequality Gap  
Narrowed

### Largest Inequality Gaps

SDR Drug Related (90%)	Teenage Birth Rate (U20) (65%)	SAR Self-Harm (61%)	SAR Alcohol Related (60%)	SAR Drug Related (52%)
------------------------	--------------------------------	---------------------	---------------------------	------------------------

### Changes in Inequality Gaps

Most Notable <b>Narrowed</b> Deprivation Inequality Gaps	Most Notable <b>Widened</b> Deprivation Inequality Gaps
Crude Intentional Self-Harm Rate U75 Cancer Mortality Potential Years Life Lost	Male Life Expectancy at 65 Alcohol Related Admissions Emergency Admissions

This is a summary of findings only. For a full assessment and all figures see downloadable tables at: <https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

## APPENDICES

### APPENDIX A: SOCIAL GRADIENT OF HEALTH

Health inequalities are often considered in terms of the gap between the most and least deprived quintiles of the population. However this does not account for those areas of intermediate levels of deprivation that may also be relatively disadvantaged in terms of their health status. The Marmot Review<sup>24</sup> demonstrated that there is a social gradient in health that runs from top to bottom of the socioeconomic spectrum, meaning that health inequalities affect everyone. There is consistent evidence from throughout the world that people at a socioeconomic disadvantage suffer a heavier burden of illness and have higher mortality rates than their better off counterparts.

Different inequality measures can give information about different aspects of inequalities. Some measures concentrate on the extremes of deprivation such as the most-least deprived (*or absolute*) gap analysis presented in the main body of this report, whilst others include relative inequality gaps across the socioeconomic scale – taking into account the whole population - and can give quite different interpretations of inequalities. Therefore, in addition to the most-least deprived (*or absolute*) gap analysis presented in this report, a social gradient analysis using the Relative Index of Inequalities (RII) has been undertaken to provide a fuller assessment of inequalities.

**Absolute gap (most-least deprived gap):** This measure describes the absolute difference between the extremes of deprivation. It has the advantage that it is intuitive and straightforward to explain, but the disadvantage that, because it focuses only on the extremes of deprivation, it does not take account of patterns of inequalities observed across the intermediate groups.

**Slope Index of Inequality (SII):** SII describes the gradient of health observed across the deprivation scale. While the absolute gap shows the difference between two large groups, SII measures the difference in health outcomes between the theoretical most and least deprived individuals, according to linear regression across health outcomes for all deprivation deciles. SII therefore has the advantage of being sensitive to the experience of the entire population, rather than just the extremes of deprivation.

**Relative Index of Inequality (RII):** The RII describes the gradient of health observed across the deprivation scale, relative to the average for the observed population (by dividing the Slope of Index of Inequality (SII) by the mean). The value of RII tells you the magnitude of inequality in relation to the mean thus representing the proportionate change in the health outcome across the population. It allows inequalities to be compared and contrasted across a number of different health indicators, and also to be monitored over time.

For further information regarding the RII methodology, including how it is calculated, please refer to the NI Health & Social Care Inequalities Monitoring System – Regional 2014 report: <https://www.health-ni.gov.uk/publications/ni-health-and-social-care-inequalities-monitoring-system-hscims-regional-2014>

<sup>24</sup> *Fair Society, Healthy Lives: The Marmot Review* can be accessed at <http://www.instituteofhealthequity.org/resources-reports/fair-society-healthy-lives-the-marmot-review>

## RESULTS

Social gradient analyses were carried out for the majority of indicators included in the HSCIMS. For some indicators this analysis could not be performed due to limitations on the level of data available. An explanatory interpretation of RII results is provided for life expectancy at birth below, with a time series for the Absolute Gap (most-least deprived) and Relative Index of Inequality (RII) for all other indicators provided in Table 4. It should be noted that in this report the SII value tends to be larger when compared with the absolute gap. This is due to the SII calculating across deprivation deciles whereas the absolute gap is calculated across deprivation quintiles.


The symbols used in the main body of the report to show the change in the inequality gap indicated by the absolute gap analysis, have also been included in the table for each indicator.

Worked examples are shown below for SII and RII.<sup>25</sup>

### Male Life Expectancy at Birth – SII

Year	2013-15	2014-16	2015-17	2016-18	2017-19
<b>Absolute Gap (Most-Least Deprived)</b>	6.5	6.6	7.1	7.1	7.0
<b>Slope Index of Inequality (SII)</b>	7.6	7.8	8.2	8.1	8.0


For life expectancy values, the Slope Index of Inequality (SII) indicates the absolute gap across the deprivation scale, represented by the gradient of the linear best fit line shown. In 2017-19, male life expectancy at birth indicates a SII gap of 8.0 years. This is higher than that indicated by the absolute gap between the most and least deprived quintiles (7.0 years). Both the absolute gap and SII showed no change in the male life expectancy deprivation gap between 2013-15 and 2017-19.



### SDR – All Cause Mortality (U75) - RII

Year	2011-15	2012-16	2013-17	2014-18	2015-19
<b>Absolute Gap (Most-Least Deprived)</b>	113%	114%	116%	115%	120%
<b>Relative Index of Inequality (RII)</b>	-0.93	-0.95	-0.95	-0.95	-0.96

The Relative Index of Inequality (RII) indicates the relative gap across the deprivation scale. In 2015-19, mortality rates among those aged below 75 years indicated a deprivation gap of -0.96, meaning that the SII value of -350 deaths per 100,000 population is equivalent to 96% of the average mortality rate across NI. Both the RII and the absolute gap showed that the deprivation gap remained similar across the analysed period.



<sup>25</sup> Calculation is deprivation quintile based i.e. difference between 20% most and least deprived areas.

**Table 4: Social Gradient Analysis of Indicators**

A comparison of the inequality gaps provided in the main body of the report with social gradient results are presented in the table below:

Indicator		Time Series				
Male Life Expectancy at Birth		2013-15	2014-16	2015-17	2016-18	2017-19
Absolute Gap (Most–Least Deprived)	—	6.5	6.6	7.1	7.1	7.0
Slope Index of Inequality (SII)	—	7.6	7.8	8.2	8.1	8.0
Female Life Expectancy at Birth		2013-15	2014-16	2015-17	2016-18	2017-19
Absolute Gap (Most–Least Deprived)	—	4.5	4.5	4.5	4.4	4.8
Slope Index of Inequality (SII)	—	5.1	5.2	5.3	5.1	5.5
Male Life Expectancy at Age 65		2013-15	2014-16	2015-17	2016-18	2017-19
Absolute Gap (Most–Least Deprived)	—	2.6	2.9	3.0	3.2	3.0
Slope Index of Inequality (SII)	—	3.1	3.3	3.3	3.5	3.3
Female Life Expectancy at Age 65		2013-15	2014-16	2015-17	2016-18	2017-19
Absolute Gap (Most–Least Deprived)	—	2.4	2.4	2.3	2.4	2.5
Slope Index of Inequality (SII)	—	2.6	2.7	2.6	2.6	2.7
Potential Years of Life Lost		2013-15	2014-16	2015-17	2016-18	2017-19
Absolute Gap (Most–Least Deprived)	◀▶	119%	118%	131%	127%	130%
Relative Index of Inequality (RII)	—	-0.98	-0.98	-1.01	-0.99	-1.01
SDR – Treatable		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	▶▶	104%	101%	103%	90%	92%
Relative Index of Inequality (RII)	▶▶	-0.85	-0.82	-0.85	-0.78	-0.78
SDR – Preventable		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	◀▶	175%	178%	177%	184%	195%
Relative Index of Inequality (RII)	—	-1.27	-1.28	-1.27	-1.28	-1.31
SDR – Avoidable		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	—	148%	149%	150%	148%	156%
Relative Index of Inequality (RII)	—	-1.12	-1.13	-1.13	-1.12	-1.14
SDR – Avoidable Children & Young People		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	—	72%	56%	56%	48%	77%
Relative Index of Inequality (RII)	—	-0.64	-0.58	-0.64	-0.59	-0.72
SDR - Circulatory (U75)		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	—	146%	143%	141%	136%	144%
Relative Index of Inequality (RII)	—	-1.10	-1.08	-1.06	-1.01	-1.04
SDR - Respiratory (U75)		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	—	228%	242%	264%	258%	249%
Relative Index of Inequality (RII)	—	-1.43	-1.47	-1.49	-1.53	-1.52
SDR - Cancer (U75)		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	—	72%	73%	72%	70%	72%
Relative Index of Inequality (RII)	—	-0.70	-0.72	-0.70	-0.68	-0.66
SDR - All Cause Mortality (U75)		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	—	113%	114%	116%	115%	120%
Relative Index of Inequality (RII)	—	-0.93	-0.95	-0.95	-0.95	-0.96
SAR - Circulatory		2013/14-15/16	2014/15-16/17	2015/16-17/18	2016/17-18/19	2017/18-19/20
Absolute Gap (Most–Least Deprived)	▶▶	27%	27%	26%	24%	22%
Relative Index of Inequality (RII)	▶▶	-0.27	-0.27	-0.25	-0.24	-0.22
SAR - Circulatory (U75)		2013/14-15/16	2014/15-16/17	2015/16-17/18	2016/17-18/19	2017/18-19/20
Absolute Gap (Most–Least Deprived)	▶▶	39%	39%	37%	35%	34%
Relative Index of Inequality (RII)	▶▶	-0.40	-0.40	-0.39	-0.37	-0.36
SPR - Antihypertensive		2015	2016	2017	2018	2019
Absolute Gap (Most–Least Deprived)	—	21%	22%	24%	23%	21%
Relative Index of Inequality (RII)	—	-0.23	-0.24	-0.25	-0.25	-0.23
SPR - Statin		2015	2016	2017	2018	2019
Absolute Gap (Most–Least Deprived)	◀▶	27%	29%	31%	33%	34%
Relative Index of Inequality (RII)	◀▶	-0.30	-0.32	-0.34	-0.35	-0.36
SAR - Respiratory		2013/14-15/16	2014/15-16/17	2015/16-17/18	2016/17-18/19	2017/18-19/20
Absolute Gap (Most–Least Deprived)	—	95%	96%	93%	94%	95%
Relative Index of Inequality (RII)	—	-0.78	-0.78	-0.75	-0.75	-0.76
SAR - Respiratory (U75)		2013/14-15/16	2014/15-16/17	2015/16-17/18	2016/17-18/19	2017/18-19/20
Absolute Gap (Most–Least Deprived)	—	115%	118%	113%	113%	112%
Relative Index of Inequality (RII)	—	-0.91	-0.91	-0.88	-0.89	-0.88
SIR - Cancer		2008-14	2009-15	2010-16	2011-17	2012-18
Absolute Gap (Most–Least Deprived)	▶▶	27%	25%	24%	24%	22%
Relative Index of Inequality (RII)	▶▶	-0.28	-0.25	-0.25	-0.26	-0.23

SAR - All Admissions		2015/16	2016/17	2017/18	2018/19	2019/20
Absolute Gap (Most–Least Deprived)	▶▶	42%	40%	39%	39%	35%
Relative Index of Inequality (RII)	▶▶	-0.42	-0.39	-0.38	-0.39	-0.37
SAR - Emergency Admissions		2015/16	2016/17	2017/18	2018/19	2019/20
Absolute Gap (Most–Least Deprived)	▶▶	73%	73%	64%	64%	60%
Relative Index of Inequality (RII)	▶▶	-0.65	-0.63	-0.57	-0.58	-0.56
SAtr – Emergency Care Attendances		2015/16	2016/17	2017/18	2018/19	2019/20
Absolute Gap (Most–Least Deprived)	—	59%	57%	58%	57%	55%
Relative Index of Inequality (RII)	—	-0.53	-0.52	-0.53	-0.52	-0.50
SAR - Elective Inpatient Admissions		2015/16	2016/17	2017/18	2018/19	2019/20
Absolute Gap (Most–Least Deprived)	—	31%	22%	26%	36%	28%
Relative Index of Inequality (RII)	—	-0.32	-0.23	-0.26	-0.36	-0.30
SAR - Day Case Admissions		2015/16	2016/17	2017/18	2018/19	2019/20
Absolute Gap (Most–Least Deprived)	—	23%	23%	24%	23%	19%
Relative Index of Inequality (RII)	—	-0.24	-0.24	-0.26	-0.26	-0.22
SAR – Self-Harm Admissions		2011/12-15/16	2012/13-16/17	2013/14-17/18	2014/15-18/19	2015/16-19/20
Absolute Gap (Most–Least Deprived)	▶▶	276%	255%	251%	239%	215%
Relative Index of Inequality (RII)	▶▶	-1.65	-1.59	-1.56	-1.51	-1.43
Crude Int. Self-Harm Rate		2013-15	2014-16	2015-17	2016-18	2017-19
Absolute Gap (Most–Least Deprived)	▶▶	161%	151%	174%	174%	105%
Relative Index of Inequality (RII)	—	-1.00	-1.09	-1.17	-1.18	-0.92
SPR - Mood & Anxiety		2015	2016	2017	2018	2019
Absolute Gap (Most–Least Deprived)	◀▶	62%	65%	67%	67%	66%
Relative Index of Inequality (RII)	◀▶	-0.58	-0.60	-0.62	-0.62	-0.62
SAR - Alcohol Related Causes		2013/14-15/16	2014/15-16/17	2015/16-17/18	2016/17-18/19	2017/18-19/20
Absolute Gap (Most–Least Deprived)	▶▶	377%	363%	338%	300%	277%
Relative Index of Inequality (RII)	▶▶	-2.0	-2.0	-1.9	-1.8	-1.7
SDR - Alcohol Specific		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	▶▶	348%	336%	353%	316%	319%
Relative Index of Inequality (RII)	—	-1.84	-1.87	-1.85	-1.83	-1.84
SDR - Smoking Related Causes		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	—	126%	129%	122%	125%	128%
Relative Index of Inequality (RII)	—	-0.98	-1.00	-0.97	-0.99	-0.99
SIR - Lung Cancer		2008-14	2009-15	2010-16	2011-17	2012-18
Absolute Gap (Most–Least Deprived)	—	165%	154%	156%	168%	164%
Relative Index of Inequality (RII)	—	-1.18	-1.12	-1.16	-1.20	-1.18
SDR - Lung Cancer		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	—	164%	163%	154%	161%	159%
Relative Index of Inequality (RII)	—	-1.20	-1.20	-1.19	-1.22	-1.18
SAR - Drug Related Causes		2013/14-15/16	2014/15-16/17	2015/16-17/18	2016/17-18/19	2017/18-19/20
Absolute Gap (Most–Least Deprived)	▶▶	267%	276%	282%	260%	239%
Relative Index of Inequality (RII)	▶▶	-1.64	-1.66	-1.66	-1.60	-1.50
SDR - Drug Related Causes		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	◀▶	347%	397%	334%	391%	404%
Relative Index of Inequality (RII)	◀▶	-1.92	-2.00	-2.00	-2.13	-2.10
SDR - Drug Misuse		2011-15	2012-16	2013-17	2014-18	2015-19
Absolute Gap (Most–Least Deprived)	◀▶	398%	464%	393%	420%	456%
Relative Index of Inequality (RII)	◀▶	-1.99	-2.08	-2.06	-2.12	-2.14
Smoking During Pregnancy		2015	2016	2017	2018	2019
Absolute Gap (Most–Least Deprived)	◀▶	305%	353%	376%	457%	367%
Relative Index of Inequality (RII)	◀▶	-1.57	-1.71	-1.75	-1.79	-1.71
Teenage Birth Rate (U20)		2015	2016	2017	2018	2019
Absolute Gap (Most–Least Deprived)	—	475%	485%	310%	400%	519%
Relative Index of Inequality (RII)	—	-1.90	-2.06	-1.67	-1.95	-1.99
Breastfeeding on Discharge		2015	2016	2017	2018	2019
Absolute Gap (Most–Least Deprived)	▶▶	50%	50%	48%	48%	46%
Relative Index of Inequality (RII)	▶▶	0.78	0.78	0.73	0.73	0.73
Low Birth Weight		2015	2016	2017	2018	2019
Absolute Gap (Most–Least Deprived)	◀▶	17%	29%	32%	38%	44%
Relative Index of Inequality (RII)	◀▶	-0.27	-0.26	-0.33	-0.42	-0.53
Small for Gestational Age <sup>26</sup>			2016	2017	2018	2019
Absolute Gap (Most–Least Deprived)			79%	43%	75%	78%
Relative Index of Inequality (RII)			-0.65	-0.51	-0.69	-0.74



Primary 1 BMI: Obese		2013/14-15/16	2014/15-16/17	2015/16-17/18	2016/17-18/19	2017/18-19/20
Absolute Gap (Most–Least Deprived)	—	45%	44%	45%	60%	64%
Relative Index of Inequality (RII)	—	-0.46	-0.47	-0.43	-0.50	-0.56
Primary 1 BMI: Overweight or Obese		2013/14-15/16	2014/15-16/17	2015/16-17/18	2016/17-18/19	2017/18-19/20
Absolute Gap (Most–Least Deprived)	—	24%	22%	21%	31%	32%
Relative Index of Inequality (RII)	—	-1.00	-0.93	-0.89	-1.17	-1.42
Year 8 BMI: Obese <sup>27</sup>		2015/16	2016/17	2017/18	2018/19	2019/20
Absolute Gap (Most–Least Deprived)		98%	107%	137%	128%	127%
Relative Index of Inequality (RII)		-0.78	-0.76	-1.18	-0.94	-0.79
Year 8 BMI: Overweight or Obese		2015/16	2016/17	2017/18	2018/19	2019/20
Absolute Gap (Most–Least Deprived)		42%	56%	55%	54%	52%
Relative Index of Inequality (RII)		-1.94	-1.97	-1.35	-1.52	-1.62

### Changes in Inequality Gaps

In the majority of indicators, there was an agreement in the assessment of change between the absolute gap and the slope index of inequality or relative index of inequality. When differences were observed, it was the result of a change in the absolute gap with the social gradient showing the gap remained constant.

The absolute deprivation gap of the following indicators widened while the social gradient analysis showed that the inequality remained constant:

- Potential Years of Life Lost
- SDR – Preventable

The absolute deprivation gap of the following indicators narrowed while the social gradient analysis showed that the inequality remained constant:

- CDR – Intentional Self-Harm
- SDR – Alcohol Specific

<sup>26</sup> No assessment of the inequality gap trend will be made until 5 years of data are available.

<sup>27</sup> No assessment of Year 8 Obesity will be made as data for the Western Trust since 2018/19 is unavailable. A full explanation of any data issues can be found in [Appendix E: Technical Notes & Definitions](#).



## Ranking of Inequality Gaps

The table below displays, in rank order from largest to smallest, the ten indicators with the largest inequality gaps as identified by RII and absolute gap analysis. As can be seen, nine of the ten indicators identified in each analysis were the same, with a few differences in the rank order of these inequality gaps.

Rank	Absolute Gap	RII	Rank Change
1	Teenage Birth Rate (U20)	SDR – Drug Misuse	1 ^
2	SDR – Drug Misuse	SDR – Drug Related Causes	1 ^
3	SDR – Drug Related Causes	Teenage Birth Rate (U20)	2 v
4	Smoking During Pregnancy	SDR – Alcohol Specific	1 ^
5	SDR – Alcohol Specific	SAR – Alcohol Related Causes	1 ^
6	SAR – Alcohol Related Causes	Smoking During Pregnancy	2 v
7	SDR – Respiratory (U75)	SDR – Respiratory (U75)	
8	SAR – Drug Related Causes	SAR – Drug Related Causes	
9	SAR – Self Harm Admissions	SAR – Self-Harm Admissions	
10	SDR – Preventable	Primary 1 BMI: Overweight or Obese	

It should be noted that life expectancy gaps have not been included in the ranking of inequality gaps above. This is because proportionately, life expectancy gaps are comparatively lower to those ranking highest in the table above. However, as the gap refers to years of life, and as life expectancy is an overarching indicator of health status it is a vital statistic of high importance and reducing this gap is considered a high priority.<sup>28</sup>


<sup>28</sup> Life expectancy inequality gaps are included as key overarching indicators of the public health strategic framework 'Making Life Better' [www.health-ni.gov.uk/articles/making-life-better-strategic-framework-public-health](http://www.health-ni.gov.uk/articles/making-life-better-strategic-framework-public-health)

### APPENDIX B: POPULATION ATTRIBUTABLE RISK (PAR) OF DEPRIVATION

Population Attributable Risk (PAR) measures the proportion of a disease/outcome (i.e. prevalence, mortality, admissions etc.) in the population that is attributable to deprivation and thus could be eliminated if deprivation were eliminated. This allows us to determine the proportional decrease in alcohol-related admissions in the population for example, in the hypothetical situation that all individuals had the same rate of alcohol-related admission as those in the highest socioeconomic category (least deprived deprivation decile). The PAR is calculated as the rate of disease in the overall population minus the rate in the unexposed group (least deprived).

PAR has been calculated in the table below for a number of health outcomes. As can be seen, the PAR percentage for preventable mortality in 2015-19 was 45% which indicates that almost half of preventable deaths in Northern Ireland were attributable to deprivation. This is reflected in the chart below showing the standardised death rate for preventable mortality by deprivation decile, with the proportion of deaths above the red line totalling the 45% of preventable deaths attributable to deprivation.

Indicator	%PAR
Teenage Birth Rate (U20)	75%
SDR - Alcohol Related Causes (U75)	54%
SAR - Self Harm Admissions	53%
SAR - Alcohol Related Causes	50%
SDR - Respiratory (U75)	49%
SDR - Preventable	45%
SIR – Lung Cancer	41%
SDR – Circulatory (U75)	40%
SDR - Avoidable	35%
Crude Death Rate - Intentional Self-harm	30%
SAR – Emergency	26%
SDR - Cancer (U75)	24%



## APPENDIX C: ADDITIONAL INDICATORS

The tables below refer to additional indicators which form part of the HSCIMS that have not been included in the main body of the report. For each indicator the figures are presented for NI, the 20% most deprived areas, the 20% least deprived areas and the most-least deprived inequality gap. In addition the RII is provided, where appropriate.

Median Fire Response Times <sup>29</sup> Time (Minutes: Seconds)	2015/16 All	2016/17 All	2017/18 All	2018/19 All	2019/20 All
<b>NI</b>	00:07:49	00:08:02	00:08:11	00:08:23	00:08:15
<b>Most Deprived</b>	00:07:17	00:06:30	00:06:38	00:06:40	00:06:42
<b>Least Deprived</b>	00:07:59	00:07:58	00:08:07	00:08:13	00:08:09
<b>Most-Least Deprived</b>	-9%	-18%	-18%	-19%	-18%

Median Ambulance Response Times <sup>29,30</sup> Time (Minutes: Seconds)	2016 All	2017 All	2018 All	2019 All	2020 All
<b>NI</b>	00:09:57	00:10:36	00:13:36	00:15:36	00:17:14
<b>Most Deprived</b>	00:07:46	00:07:38	00:09:41	00:12:11	00:14:26
<b>Least Deprived</b>	00:11:02	00:12:58	00:14:48	00:16:21	00:17:51
<b>Most-Least Deprived</b>	-30%	-41%	-20%	-25%	-19%

SDR - All Age All Cause Mortality Deaths per 100,000 population	2011-15 All	2012-16 All	2013-17 All	2014-18 All	2015-19 All
<b>NI</b>	1,041	1,036	1,032	1,023	1,016
<b>Most Deprived</b>	1,252	1,250	1,241	1,246	1,253
<b>Least Deprived</b>	908	900	901	887	879
<b>Most-Least Deprived</b>	38%	39%	38%	40%	43%
<b>RII</b>	-0.37	-0.39	-0.38	-0.40	-0.42

Looked After Children Rate per 1,000 population under 18 years	2014 All	2015 All	2016 All	2017 All	2018 All
<b>NI</b>	5.0	5.0	5.1	5.3	5.5
<b>Most Deprived</b>	10.0	10.0	10.5	10.8	11.3
<b>Least Deprived</b>	1.7	1.7	1.5	1.6	1.8
<b>Most-Least Deprived</b>	484%	482%	582%	555%	530%

Autism Prevalence in School Age Children Rate per 100,000 population	2015/16 All	2016/17 All	2017/18 All	2018/19 All	2019/20 All
<b>NI</b>	2,310	2,509	2,909	3,331	4,237
<b>Most Deprived</b>	2,844	3,207	3,598	4,321	5,515
<b>Least Deprived</b>	2,277	2,332	2,861	3,223	4,154
<b>Most-Least Deprived</b>	25%	37%	26%	34%	33%

<sup>29</sup> Evidence shows that emergency times are correlated more with location such as urban/rural than deprivation.

<sup>30</sup> In 2019/20, a new Clinical Response Model (CRM) programme was introduced along with a new set of ambulance categories in line with the national Ambulance Response Programme (ARP). Therefore, information from 2019/20 onwards is calculated differently and cannot be directly compared with previous years.

**Changes to Avoidable Mortality Definitions:** following an Office for National Statistics (ONS) consultation,<sup>31</sup> on a new definition of avoidable mortality as proposed by the Organisation for Economic Co-operation and Development (OECD),<sup>32</sup> a new definition was implemented in 2020. Figures based on the old definition remain included below for Programme for Government (PfG) monitoring purposes, specifically the Preventable Mortality indicator. Full details can be found in [Appendix E: Technical Notes & Definitions](#). The tables below provide figures based on the previous ONS definition, the full definition of which can be found in Table 13 within Appendix E.

SDR Amenable Mortality (Previous ONS Definition) Rate per 100,000 population	2011/15	2012/16	2013/17	2014/18	2015/19
	All	All	All	All	All
<b>NI</b>	127	127	124	123	139
<b>Most Deprived</b>	201	199	196	193	217
<b>Least Deprived</b>	89	88	86	87	90
<b>Most-Least Deprived</b>	127%	127%	128%	121%	142%

SDR Preventable Mortality (Previous ONS Definition) Rate per 100,000 population	2011/15	2012/16	2013/17	2014/18	2015/19
	All	All	All	All	All
<b>NI</b>	207	205	207	207	208
<b>Most Deprived</b>	338	335	335	333	336
<b>Least Deprived</b>	139	139	142	141	141
<b>Most-Least Deprived</b>	144%	141%	135%	136%	138%

SDR Avoidable Mortality (Previous ONS Definition) Rate per 100,000 population	2011/15	2012/16	2013/17	2014/18	2015/19
	All	All	All	All	All
<b>NI</b>	245	242	244	244	244
<b>Most Deprived</b>	398	391	390	388	391
<b>Least Deprived</b>	167	166	168	169	167
<b>Most-Least Deprived</b>	138%	136%	132%	130%	135%

SDR Avoidable: Children & Young People Mortality (Previous ONS Definition) Rate per 100,000 population	2011/15	2012/16	2013/17	2014/18	2015/19
	All	All	All	All	All
<b>NI</b>	24	22	22	22	22
<b>Most Deprived</b>	33	29	28	28	28
<b>Least Deprived</b>	20	19	18	19	16
<b>Most-Least Deprived</b>	68%	53%	54%	46%	76%

<sup>31</sup> <https://consultations.ons.gov.uk/health-and-life-events/avoidable-mortality-definition/>

<sup>32</sup> <http://www.oecd.org/health/health-systems/Avoidable-mortality-2019-Joint-OECD-Eurostat-List-preventable-treatable-causes-of-death.pdf>

## APPENDIX D: URBAN-RURAL ANALYSIS

Urban-Rural analysis included below is based on the 2015 NISRA Urban-Rural classification, with the exception of Healthy Life Expectancy and Disability Free Life Expectancy which use the 2005 urban rural classification for all years up to and including 2015, due to data limitations. Further information regarding urban-rural classification can be found on the NISRA webpage at <https://www.nisra.gov.uk/support/geography/urban-rural-classification>.

**A positive inequality gap means that the health outcomes in Northern Ireland are worse than in the rural areas.**

### Summary of findings

Compared with the regional average, rural areas experienced better outcomes across the majority of indicators analysed, however fire and ambulance response times continue to remain higher in rural areas.

### Outcomes that were significantly better in rural areas than the NI average

Male Life Expectancy at Birth	SDR Cancer (U75)	SPR Mood & Anxiety
Female Life Expectancy at Birth	SDR All Cause (U75)	SAR Alcohol
Male Healthy Life Expectancy	SPR Antihypertensive	SDR Alcohol
Female Healthy Life Expectancy	SPR Statin	SDR Smoking
Male Disability Free Life Expectancy	SAR Respiratory	SIR Lung Cancer
Female Disability Free Life Expectancy	SAR Respiratory (U75)	SDR Lung Cancer
Male Life Expectancy at 65	SAR Circulatory	SAR Drug Related
Female Life Expectancy at 65	SIR Cancer	SDR Drug Related
PYLL	SAR All	SDR Drug Misuse
SDR Treatable	SAR Emergency	Infant Mortality
SDR Preventable	SAtR Emergency Care	Smoking During Pregnancy
SDR Avoidable	SAR Day Case	Teenage Birth Rate
SDR Circulatory (U75)	SAR Self Harm	Small for Gestational Age
SDR Respiratory (U75)	CDR Intentional Self-Harm	P1 Obese

### Outcomes that were significantly worse in rural areas than the NI average

Ambulance Response Times

Fire Response Times

### Outcomes that were similar (or not significantly different) in Rural areas and the NI average

SDR Avoidable: Children & Young People	SAR Elective Inpatient Admissions	Low Birth Weight
SAR Circulatory (U75)	Healthy Birth Weight	Breastfeeding on Discharge
		P1 Overweight or Obese

Figures for each indicator for NI, Rural areas, Urban areas, Mixed Urban-Rural areas and the NI-Rural Gap, are provided within the accompanying downloadable tables:

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>

## APPENDIX E: TECHNICAL NOTES & DEFINITIONS

### Indicators

There are 51 indicators included in the Northern Ireland analyses for the current report. Not all indicators are assessed at each level of geography (see [Table 5](#)), and dependent on the number of years data available, or any potential quality issues, assessments may not be made on all aspects of an indicator.

Due to random fluctuations in events over time, it is often necessary to aggregate more than one year of data for indicators, in order to ensure stability. The number of years of information that are required to aggregate for each indicator is informed by both the number of events and also an assessment of its annual variability.

### Standardisation Methods

A number of indicators included in this report have been age standardised to allow the comparison of rates between populations with different age structures by relating them to a standard population, in this case the 2013 European Standard Population (90+ version). In most circumstances direct standardisation is used which not only allows the comparison of disease and death rates across both areas and time, but also to assess the relative burden of disease in a population.

### Indicator Stability/Confidence Intervals

Indicator stability at the regional level does not mean that an indicator is also stable at the lower geographic levels of HSC Trust, LGD or DEA. To ensure robustness of the data, confidence intervals were calculated for rates for the most recent year at each geographic level, including the 20% most deprived Trust and LGD areas. The confidence interval for each standardised rate was assessed, in terms of its size and in relation to other comparable rates for other geographical areas, i.e. the Belfast Trust average and its 20% most deprived Trust areas. As a result of these assessments not all of the 51 indicators examined at the regional level were deemed robust enough to be presented at the sub-regional level, of these 51 indicators; 45 were found suitable to be published at the HSC Trust level, 43 at the LGD level and 30 at the DEA level. A full list of indicators and the level analysed can be found in [Table 5](#).

Confidence intervals are used to quantify the imprecision in the estimate of a particular value. Specifically it quantifies the imprecision that results from random variation in the estimation of the value. In many cases the source of this random variation is sampling, for example in Healthy Life Expectancy, as any measurement taken from a sample provides an imprecise estimate of the true population value. In public health many indicators are based on what can be considered to be complete data sets and not samples, e.g. age standardised mortality rates based on death registers. In these instances the imprecision arises not as a result of sampling variation but of 'natural' variation. The indicator is considered to be the outcome of a stochastic process, i.e. one which can be influenced by the random occurrences that are inherent in the world around us. In such instances the value actually observed is only one of the set that could occur under the same circumstances. Generally in public health, it is the underlying circumstances or process that is of interest and the actual value observed gives only an imprecise estimate of this 'underlying risk'.

## Review of Suicide Statistics in Northern Ireland

Suicide deaths in Northern Ireland are defined as deaths from Self-inflicted Injury (also referred to as intentional self-harm) as well as Events of Undetermined Intent. This is consistent with the UK National Statistics definition. A death which is suspected to be suicide must be referred to the Coroner with the information provided by coroners at registration of the death is used to code the underlying cause of death. In some instances, it can be difficult to establish whether the cause of death was suicide. If it is not clear, or the Coroner has not specifically stated that it is a suicide, these are coded as 'Undetermined'. Following a quality exercise between NISRA Vital Statistics Unit and the Coroners' Service, to better understand drug related deaths and intent, improvements have been made in order to reduce the number of deaths coded as 'undetermined' since Quarter 3 2018. These changes have resulted in a discontinuity in the statistical series, with the number of 'undetermined' deaths reducing from 132 in 2017 to 10 in 2019. NISRA is working with the Coroners' Service to review and revise, as necessary, drug related deaths within the 'undetermined deaths' category from 2015 to 2018. NISRA recommends that until this review has completed users should refer to the sub-series relating to self-inflicted injury (also referred to as intentional self-harm) only, as this is unaffected by the discontinuity outlined above and remains a reliable indication of the trend in suicides over recent years. Further information on this review and detailed statistics on the number of suicides registered each year in Northern Ireland can be accessed at the link below.

<https://www.nisra.gov.uk/publications/suicide-statistics>

## Use of Suicide Statistics in this Publication

In previous iterations of this report, and other reports produced by PHIRB, mortality from suicide is calculated according to the UK National Statistics definition shown in the table below.

ICD-10 Underlying Cause Code	Description
X60-84, Y87.0	Self-inflicted Injury (Intentional self-harm)
Y10-Y34, Y87.2	Events of Undetermined Intent

In line with the review and advice from Vital Statistics Unit this definition has not been used in this publication. The sub-series relating to self-inflicted injury/intentional self-harm will be used in its place in order to ensure comparability across years. As such, the crude death rate indicator for suicide has been replaced by a crude deaths rate indicator for intentional self-harm. It is envisaged that following the completion of the review subsequent editions of this publication will revert to original UK National Statistics definition for suicide.

Table 5: HSCIMS Indicators Analysed at Northern Ireland (NI), HSC Trust, LGD and DEA level

INDICATOR	NI	Trust	LGD	DEA
Male Life Expectancy at Birth	●	●	●	●
Female Life Expectancy at Birth	●	●	●	●
Male Life Expectancy at Age 65	●	●	●	●
Female Life Expectancy at Age 65	●	●	●	●
Male Healthy Life Expectancy	●			
Female Healthy Life Expectancy	●			
Male Disability Free Life Expectancy	●			
Female Disability Free Life Expectancy	●			
Potential Years of Life Lost –All	●	●	●	●
Standardised Death Rate – Treatable	●	●	●	●
Standardised Death Rate – Preventable	●	●	●	●
Standardised Death Rate – Avoidable	●	●	●	●
Standardised Death Rate – Avoidable: Children & Young People	●			
Standardised Death Rate - Circulatory (U75)	●	●	●	
Standardised Death Rate - Respiratory (U75)	●	●		
Standardised Death Rate - Cancer (U75)	●	●	●	
Standardised Death Rate - All Cause Mortality (U75)	●	●	●	
Standardised Admission Rate –Circulatory	●	●	●	●
Standardised Admission Rate - Circulatory (U75)	●	●	●	●
Standardised Prescription Rate – Antihypertensive	●	●	●	●
Standardised Prescription Rate – Statin	●	●	●	●
Standardised Admission Rate – Respiratory	●	●	●	●
Standardised Admission Rate - Respiratory (U75)	●	●	●	●
Standardised Incidence Rate – Cancer	●	●	●	●
Standardised Admission Rate - All Admissions	●	●	●	●
Standardised Admission Rate - Emergency Admissions	●	●	●	●
Standardised Attendance Rate - Emergency Care	●	●	●	
Standardised Admission Rate - Elective Inpatient Admissions	●	●	●	●
Standardised Admission Rate - Day Case Admissions	●	●	●	●
Standardised Admission Rate – Self-Harm Admissions	●	●	●	
Crude Suicide Rate	●	●	●	●
Standardised Prescription Rate - Mood & Anxiety	●	●	●	●
Standardised Admission Rate - Alcohol Related Causes	●	●	●	●
Standardised Death Rate - Alcohol Specific Causes	●	●	●	
Standardised Death Rate - Smoking Related Causes	●	●	●	●
Standardised Incidence Rate - Lung Cancer	●	●	●	●
Standardised Death Rate - Lung Cancer	●	●	●	
Standardised Admission Rate - Drug Related Causes	●	●	●	●
Standardised Death Rate - Drug Related Causes	●	●	●	
Standardised Death Rate - Drug Misuse	●	●		
Infant Mortality Rate	●			
Smoking During Pregnancy	●	●	●	●
Teenage Birth Rate (U20)	●	●	●	
Low Birth Weight	●	●	●	●
Healthy Birth Weight	●	●	●	●
Breastfeeding on Discharge	●	●	●	●
Small for Gestational Age	●	●	●	●
Primary 1 BMI: Obese	●	●	●	
Primary 1 BMI: Obese & Overweight	●	●	●	
Year 8 BMI: Obese	●	●	●	
Year 8 BMI: Obese & Overweight or Obese	●	●	●	



## Methodology for Assessing Health Outcomes

In order to provide an assessment of the LGD to NI inequality gaps for the most recent year, analysis was performed to indicate whether the LGD average was better than, similar to, or worse than the NI average. If the LGD average of the health outcome had overlapping confidence intervals with the NI average, then the health outcome was reported as being similar to the NI average. Where confidence intervals did not overlap, the LGD average of the health outcome was reported as being either better or worse than the NI average. This methodology was employed for all standardised rates (i.e. death, admission, incidence, and prescription rates). For those health outcomes which did not have confidence intervals associated with them, such as teenage birth rate, a range of +/- 5% was calculated for each health outcome value and if the NI average fell within this range the health outcome was considered to be similar to the NI average. It should be noted that given the particular sensitivity around the health outcome 'crude intentional self-harm rate' and the relatively small numbers involved a range of +/- 2.5% was employed. This methodology allowed us to identify any health outcomes which were notably worse or better than the NI average and was used to provide an assessment of the health outcomes at DEA level, compared with the LGD average. As with all of our observations of differences between areas and assessments of changes over time, conclusions are open to interpretation.

## Mortality Rates

For simplicity of understanding, mortality figures are based on the single main underlying cause of death classification, but a death can be due to a variety of different causes. This can lead to an underestimation of the impact of common conditions associated with multiple causes of death (e.g. diabetes, influenza and pneumonia). All death figures used in the HSCIMS are based on the year that the death was registered and not necessarily the year in which the death occurred. While the vast majority of deaths are registered shortly after death, there may be a delay in registering some deaths. Events such as infant death or suicide are usually referred to a coroner and this legal process can take some time.

## Population

Population is a vital part of rate calculations; a change to the size of a population or its age distribution will impact on rates and subsequently inequality gaps. For instance, overall yearly deaths in Northern Ireland remained between 14,000 and 16,250 from the turn of the century up to 2019, yet mortality rates have been falling – this can be partially explained by the growing and ageing Northern Ireland population. Between 2008 and 2019 for example, the population grew from 1,779,152 to 1,893,667; an increase of 114,515 persons (6.4%). During this time the proportion of the population aged 65 and over increased from 13.9% (247,500 persons) in 2008 to 16.6% (314,724 persons) in 2019.

## Small Area Population Estimates


Population estimates disaggregated to a relatively small geographic area level (i.e. Super Output Area (SOA) and Small Area (SA)), by age and gender, are used to calculate many of the HSCIMS indicators for deprived and rural areas. However, as population estimates produced for NI are not available to the required level of detail, it is necessary to rework these estimates by proportioning out aggregated small area population estimates by gender and single year of age breakdowns from NISRA mid-year estimates. These reworked estimates are validated by a process of integrity checks with higher level age and geography population totals published by NISRA. Reworked estimates are calculated from unrounded population breakdown figures which may not match exactly with some of the population breakdowns published by NISRA which have been rounded to the nearest person.

## Deprivation Classification

The deprivation classification used in this report is based on the Northern Ireland Multiple Deprivation Measure (NIMDM) produced by NISRA. The 20% most and least deprived areas are defined according to the NIMDM 2017.<sup>33</sup>

Although the 2017 NIMDM is available at small area level it was decided to continue using the SOA classification within the HSCIMS to ensure continuity and comparability with the back series of data and across indicators. In addition, all analysis presented is based on multiple deprivation rather than any specific deprivation domain.

**Chart 1 – 20% Most and least deprived areas in Northern Ireland according to the 2017 NIMDM**



© Crown copyright and database rights NIMA MOU210

## Drug Related Admissions and Mortality

Please note that some observations may be due to changes in drug misuse behaviours among the population. There are ongoing concerns about polydrug misuse and the misuse of prescription drugs and new psychoactive substances. It appears that a significant cohort are engaging in increasingly risky behaviours, with an acute increase in related harms.

## Childhood Obesity

The data cleansing parameters applied to the calculation of childhood obesity figures has been refined from 2017/18 onwards. Therefore figures should be treated with caution when making comparisons.

Year 8 childhood obesity data for the Western Trust since 2018/19 is unavailable and therefore figures for NI should be treated with caution. As a result, no regional trend analysis for these indicators will be made, nor will any comparison of sub-regional areas with the NI average. Sub-regional trend analysis is available for all HSC Trusts and LGDs with the exception of the Western HSC Trust and the Causeway Coast & Glens, Derry City & Strabane and Fermanagh & Omagh LGDs. In addition, Primary 1 figures combine 3 years of data, as school closures due to the COVID-19 pandemic led to a reduction in the number of recorded measurements for this year group. Data for children in Year 8 did not appear to be affected by school closures.

<sup>33</sup> <https://www.nisra.gov.uk/statistics/deprivation/northern-ireland-multiple-deprivation-measure-2017-nimdm2017>

## Sources of Information

Table 6: Indicators and Supplementary Information

Information	Source
Deaths and births	General Register Office, Vital Statistics & Administrative Research and Support Branch (VARs), NI Statistics and Research Agency (NISRA)
Hospital Admissions & Attendances	Hospital Information Branch, Information Analysis Directorate, DoH
Prescriptions	Business Services Organisation
Cancer Incidence	NI Cancer Registry
Smoking during pregnancy, breastfeeding, low birth weight, small for gestational age, healthy birth weight	NI Maternity System <sup>34</sup>
Childhood overweight/obesity	Child Health System
Fire response times	NI Fire and Rescue Service
Ambulance response times	NI Ambulance Service
NI Health Survey	Public Health Information & Research Branch, Information Analysis Directorate, DoH
Continuous Household Survey	NI Statistics and Research Agency (NISRA)
NI Small Area Population Estimates	NI Statistics and Research Agency (NISRA)
European Standard Population (ESP) 2013	Eurostat
Deprivation classification	NI Multiple Deprivation Measure 2017 (NISRA)
Urban-rural classification	NI Statistics and Research Agency (NISRA)
Looked after Children	Community Information Branch, Information Analysis Directorate, DoH
Children with Autism	Community Information Branch, Information Analysis Directorate, DoH

## Indicator Definitions

Disease Classification - The indicators making up the HSCIMS are classified using the International Classification of Disease, 10<sup>th</sup> revision (ICD-10). This is the standard diagnostic tool for epidemiology, health management and clinical purposes, including the analysis of the general health situation of population groups.

A complete listing of ICD-10 codes can be found at the following web link:

[www.who.int/classifications/apps/icd/icd10online/](http://www.who.int/classifications/apps/icd/icd10online/)

<sup>34</sup> Please note that prior to 2017, data used in the production of low birth weight statistics, as well as healthy birth weight, smoking during pregnancy and breastfeeding, were provided from each of the HSC Trust Child Health Systems (CHS). From 2017 onwards figures are produced directly from the Northern Ireland Maternity System (NIMATS) by Information & Analysis Directorate (IAD). Low birth weight data from NIMATS data is used to populate the Trust CHS so data from the two systems should be consistent. IAD have investigated historic data to ensure that previously published data obtained through CHS was wholly consistent with that held on NIMATS.

<b>LIFE EXPECTANCY</b>	
<b>Life Expectancy Estimates</b>	NISRA publish the official life expectancy estimates at NI, Local Government District and Parliamentary Constituency level. The HSCIMS publishes at further levels to allow for assessment of inequality gaps between different areas/population groups, including deprivation analysis.
<b>Life Expectancy at Birth</b>	The expected years of life at time of birth based on mortality patterns in the period in question. It is based on the average death rates over a three year period. Presented separately for males and females.
<b>Life Expectancy at Age 65</b>	The expected years of life at age 65 based on mortality patterns in the period in question. It is based on the average death rates over a three year period. Presented separately for males and females.
<b>Healthy Life Expectancy (HLE)</b>	This is the average number of years a person can expect to live in good health. HLE provides an estimate of lifetime spent in 'Very Good' or 'Good' health, calculated using respondents' perception of their own health according to the Health Survey Northern Ireland (HSNI). HLE excludes communal establishments. All urban/rural analysis prior to 2016 is based on the 2005 urban-rural classification.
<b>Disability Free Life Expectancy (DFLE)</b>	This is the average number of years a person can expect to live disability free. DFLE provides an estimate of lifetime spent free from a limiting persistent (twelve months or more) illness or disability, based upon a self-rated functional assessment of health recorded in the HSNI. DFLE excludes communal establishments. All urban/rural analysis prior to 2016 is based on the 2005 urban-rural classification. It should be noted that the health survey question used to determine longstanding illness changed from 2012/13 onward by making specific reference to mental health conditions in addition to physical. The new question is based on the UK harmonised principle for long-lasting health conditions and illness. This change may have affected responses to the question and subsequently impacted on DFLE figures. For further information contact PHIRB (details on reverse of publication).
<b>Pregnancy &amp; Early Years</b>	
<b>Teenage Birth Rate (U20)</b>	The number of births in an area to teenage mothers (i.e. Between 13 and 19 years of age) expressed per 1,000 females.
<b>Smoking during Pregnancy</b>	The proportion of all live births, where the Health and Care Number (HCN) of the mother is recorded, that were to mothers that reported smoking during pregnancy. Information is gathered at the 'booking in' appointment and therefore represents mothers at the end of the first trimester. As this indicator is self-reported, it may be subject to a degree of under-reporting.
<b>Low Birth Weight</b>	The proportion of all live births where the HCN of the mother is recorded and the birth weight of the child was less than 2,500g.
<b>Healthy Birth Weight</b>	The proportion of all live births, where the HCN of the mother is recorded, with a birth weight within a range appropriate for their gestational age and gender.
<b>Breastfeeding on Discharge</b>	The proportion of all live births, where the HCN of the mother is recorded, that were being breastfed on discharge from hospital. Figures include mothers' breastfeeding their child as well as using complementary feeding.
<b>Small for Gestational Age</b>	The proportion of all live births, where the HCN of the mother is recorded, that were small-for-gestational age (SGA). This is when an infant is born with a birth weight less than the 10th percentile, on a chart customised for maternal characteristics, for gestational age in body weight.

ADMISSIONS	
<b>Hospital Information System (HIS)</b>	Admissions data used to calculate rates are provided by the Hospital Information Branch and are extracted from the Hospital Information System (HIS). All mental health specialities have been excluded from the data. Figures are based on number of admissions and not individuals. Further information and definition on inpatient and day case activity is available at <a href="https://www.health-ni.gov.uk/articles/inpatient-and-day-case-activity">https://www.health-ni.gov.uk/articles/inpatient-and-day-case-activity</a> .
<b>Standardised Admission Rate (SAR)</b>	This is calculated by standardising (using the direct method) the average admission rate in NI (over a predefined period) due to specified ICD-10 classification codes (may also be age specific) to the 2013 European Standard Population (ESP).
Indicator Name	
<b>- All Admissions</b>	Includes all acute inpatient and day case admissions (excluding regular day and night attenders, hospital transfers and other (maternity/delivery episodes)). Deaths and discharges have been used as an approximation for admissions.
<b>- Emergency Admissions</b>	A patient for whom admission is unpredictable and at short notice because of clinical need. All non-elective acute admissions excluding maternity, other and not known.
<b>- Elective Inpatient Admissions</b>	A patient for whom the decision to admit could be separated in time from the actual admission. Does not include day cases, not to be confused with elective admissions (which include day cases)
<b>- Day Case Admissions</b>	A patient admitted electively during the course of a day with the intention of receiving care who does not require the use of a hospital bed overnight and who returns home as scheduled. If this original intention is not fulfilled and the patient stays overnight, such a patient should be counted as an inpatient and is not counted as a day case admission.
<b>- Circulatory</b>	Selected according to International Classification of Disease (ICD-10) codes I00-I99. <sup>35</sup>
<b>- Circulatory U75</b>	ICD-10 codes I00-I99, under 75 years of age.
<b>- Respiratory</b>	ICD-10 codes J00-J99.
<b>- Respiratory U75</b>	ICD-10 codes J00-J99, under 75 years of age.
<b>- Alcohol Related Causes</b>	Alcohol related causes included in Table 8.
<b>- Drug Related Causes</b>	Drug related causes included in Table 10.
<b>- Self-Harm Admissions</b>	ICD-10 codes X60-84 and Y87.0. This indicator was developed to complement the suicide information, however it does not provide a complete picture of the problem of self-harm (or parasuicide) as in many instances, self-harm does not result in an acute admission to hospital. It should be noted that there have been a range of additional infrastructure provided to support people presenting with self-harm. These programmes may be contributing to the decrease in self-harm admissions.

<sup>35</sup> For a listing and explanation of topology or site codes see: International Statistical Classification of Diseases and Related Health Problems, Tenth Revision, World Health Organisation, Geneva. Or view online at: <http://apps.who.int/classifications/icd10/browse/2019/en#/I1>

**ATTENDANCES****SYMPHONY & EEMS**

Attendance data used to calculate rates are provided by the Hospital Information Branch and are from the two administrative systems used by emergency departments in Northern Ireland (SYMPHONY & EEMS). Figures are based on number of attendances and not individuals. Further information on emergency care activity is available at <https://www.health-ni.gov.uk/articles/emergency-care-and-ambulance-statistics>.

**Indicator Name****- Emergency Care Attendances**

New and unplanned review attendances at all Emergency Departments (Types 1 - 3). Data for RVH-RAES (Eye Casualty) not included prior to 2018/19. This relates to any patient who presents without appointment to an emergency care department. This differs from an emergency admission where a patient is admitted to an acute hospital by various routes, including through an emergency care department or via a General Practitioner.

**MORTALITY****Infant Mortality Rate**

The number of infant deaths per 1,000 live births. Infant deaths refer to all deaths in the first year of life.

**Potential Years of Life Lost (PYLL)**

This is calculated by summing the deaths occurring at each age and multiplying this with the number of years a person of that age could have been expected to live. It is a summary measure of premature mortality, weighting deaths occurring at younger ages, which are, a priori, preventable. It uses the average age-specific life expectancy for each five year age band as the age to which a person in that age band might be expected to live.

**Intentional Self-Harm (CDR)**

*Not age standardised as it was found to make little or no difference whilst introducing a new confidence interval*

The number of deaths by intentional self-harm per 100,000 population ICD-10 codes X60-X84, Y87.0.

**Standardised Death Rate (SDR)**

This is calculated by directly age standardising the average death rate in NI over a given period, due to specific causes of death (selected according to ICD-10 classification) to the 2013 European Standard Population (ESP). Some death rates relate to those under the age of 75 as indicators of premature mortality for specific diseases.

**Indicator Name****- All Cause U75**

All causes, under 75 year of age

**- All Age All Cause**

All causes

**- Treatable**

Causes of death that can be mainly avoided through timely and effective health care interventions, including secondary prevention and treatment (i.e. after the onset of disease, to reduce case-fatality) – see Table 7 for full list of causes.

**- Preventable**

Causes of death that can be mainly avoided through effective public health and primary prevention interventions (i.e. before the onset of diseases/injuries, to reduce incidence) – see Table 7 for full list of causes.

**- Avoidable**

Avoidable deaths are all those defined as preventable and treatable – see Table 7 for full list of causes.

**- Avoidable: Children & Young People**

Causes of death are the same as those for overall avoidable, with all causes restricted to those aged 0-19 years. – see Table 7 for full list of causes.

**- Circulatory U75**

ICD-10 codes I00-I99, under 75 year of age.

**- Respiratory U75**

ICD-10 codes J00-J99, under 75 year of age.

**- Cancer U75**

ICD-10 codes C00-C97, under 75 year of age.

**- Lung Cancer**

ICD-10 codes C33-C34.

**- Alcohol Specific**

Alcohol Specific causes – see Table 9 for full list of causes.

- Drug Related Causes	Drug related causes – see Table 10 for full list of causes.
- Drug Misuse	Deaths related to drug misuse – see Table 11 for full list of causes.
- Smoking Related Causes	Deaths due to Smoking related causes– see Table 12 for full list of causes.

### Cancer Incidence

#### Northern Ireland Cancer Registry (NICR)

Cancer incidence numbers are extracted from the NICR's "live" database, and hence are continuously updated. As a result, an earlier extract taken at a later date may supply slightly different results. Therefore, although the overall trend will be the same, previously published data and data published elsewhere may have rates that vary slightly to what is published is here.

NICR publish official Standardised Incidence Rates (SIRs), however the HSCIMS publishes at further levels to allow for assessment of inequality gaps between different areas/population groups, including deprivation analysis.

#### Standardised Incidence Rate (SIR)

This is calculated by standardising (using the direct method) the average incidence rate in NI (over seven years) due to specified ICD-10 classification codes to the 2013 European Standard Population (ESP).

#### Indicator Name

#### ICD-10 Classification

#### - Cancer

ICD-10 codes C00-C97, excluding C44 (non-melanoma skin cancer which is quite common, in most cases easily treated and rarely fatal).

#### - Lung Cancer

ICD-10 codes C33 and C34.

### Prescriptions

#### Electronic Prescribing Eligibility System (EPES)

Prescription data is extracted from the EPES which is maintained by Business Services Organisation (BSO). The data provided covers drugs dispensed in primary care only, and includes prescriptions issued by all types of prescribers including doctors, nurses and dentists, and all those issued and dispensed by pharmacists, dispensing doctors and appliance suppliers. Drugs prescribed and dispensed in hospital cannot be captured centrally due to the use of different IT systems.

#### Standardised Prescription Rate (SPR)

This is calculated by standardising (using the direct method) the average prescription rate (over one year) in NI for people dispensed predefined prescription drugs, to the 2013 European Standard Population (ESP). Rates refer to number of persons prescribed a drug and does not include multiple prescription.

#### Indicator Name

#### British National Formulary (BNF) code

#### - Antihypertensive

Drugs included are those with a BNF code of 2.2.1, 2.4, 2.5.5.1, 2.5.5.2 and 2.6.2

#### - Statin

Drugs included are those with a BNF code of 2.12

#### - Mood & Anxiety Disorders

Drugs included are those with a BNF code of 4.1.2 and 4.3



<b>Childhood Obesity</b>	
<b>Childhood Overweight and Obese</b>	<p>Height and weight information is extracted from the Child Health System (CHS) and converted into a Body Mass Index (BMI) score for each pupil. The BMI can be categorized using International Growth Charts as determined by the International Obesity Taskforce (IOTF) which consider age and gender, allowing the identification of those who are overweight or obese. Records are analysed based on two criteria:</p> <p>Date of Exam within the Primary 1 or Year 8 school year: 01/09/XX-31/08/XX  Date of Birth for Primary 1 or Year 8 pupils: 02/07/XX - 01/07/XX  For data since 2017/18, additional data cleansing parameters have been applied.</p>
<b>Indicator Name</b>	<b>CHS Data</b>
<b>-Primary 1 BMI: Obese</b>	The proportion of children in Primary 1 classified as obese.
<b>-Primary 1 BMI: Overweight or Obese</b>	The proportion of children in Primary 1 classified as overweight or obese.
<b>-Year 8 BMI: Obese</b>	The proportion of children in Year 8 classified as obese.
<b>-Year 8 BMI: Overweight or Obese</b>	The proportion of children in Year 8 classified as overweight or obese.
<b>Additional Indicators</b>	
<b>Median Fire Response Time</b>	<p>The median response time taken by the Northern Ireland Fire and Rescue Service (NIFRS) to respond to an incident. The 'response time' is measured as the 'time of the call to NIFRS Regional Control Centre' to 'the time the 1st Appliance books in attendance' at the incident.</p> <p>Calculations are based on the time taken for NIFRS to respond to each incident within a one year time period.</p> <p>The median i.e. midpoint value is reported rather than the simple average as it is unaffected by atypically long or short response times.</p>
<b>Median Ambulance Response Time</b>	<p>The median time taken by the appropriate response vehicle to respond to an incident. Calculations are based on the time taken to respond to each incident within a one month time period (August). This data refers to CAT1, CAT1(T) and CAT2 emergency responses, excluding Healthcare Professionals (HCP) calls.</p> <p>The median i.e. midpoint value is reported rather than the simple average as it is unaffected by atypically long or short response times.</p>
<b>Looked after Children</b>	The number of looked after children per 100,000 population (under 18 years of age) by location prior to last entering care. Data was extracted from the annual OC2 Community Information Return, which includes children who have been in care continuously for twelve months or longer at 30th September.
<b>Autism Prevalence in School Age Children</b>	The number of children with Autism or Asperger Syndrome per 100,000 children in compulsory grant-aided education. Data extracted from the NI School Census.



## ICD-10 Classification Tables

The table below lists the revised ICD-10 classification codes of all causes of death considered avoidable, with indication as to which are considered treatable, preventable or both. This definition has been implemented following an Office for National Statistics (ONS) consultation,<sup>36</sup> on a new definition of avoidable mortality as proposed by the Organisation for Economic Co-operation and Development (OECD).<sup>37</sup> All avoidable mortality indicators, including the back series, are based on the new definition. The previous definition, for use with the additional indicators in [Appendix C](#), can be found in Table 13.

Table 7: Treatable, Preventable &amp; Avoidable Causes

Condition Group & Cause	ICD-10 Codes	Age	Treatable	Preventable
<b>Infectious Diseases</b>				
Intestinal Diseases	A00-A09	0-74		•
Diphtheria, Tetanus, Poliomyelitis	A35, A36, A80	0-74		•
Whooping cough	A37	0-74		•
Meningococcal infection	A39	0-74		•
Sepsis due to streptococcus pneumonia and sepsis due to haemophilus influenza	A40.3, A41.3	0-74		•
Haemophilus influenza infections	A49.2	0-74		•
Sexually transmitted infections (except HIV/AIDS)	A50-A60, A63, A64	0-74		•
Varicella	B01	0-74		•
Measles	B05	0-74		•
Rubella	B06	0-74		•
Viral Hepatitis	B15-B19	0-74		•
HIV/AIDS	B20-B24	0-74		•
Malaria	B50-B54	0-74		•
Haemophilus and pneumococcal meningitis	G00.0, G00.1	0-74		•
Tuberculosis	A15-A19, B90, J65	0-74	• (50%)	• (50%)
Scarlet fever	A38	0-74	•	
Sepsis	A40 (excl. A40.3), A41 (excl. A41.3)	0-74	•	
Cellulitis	A46, L03	0-74	•	
Legionnaires disease	A48.1	0-74	•	
Streptococcal and enterococci infection	A49.1	0-74	•	
Other meningitis	G00.2, G00.3, G00.8, G00.9	0-74	•	
Meningitis due to other and unspecified causes	G03	0-74	•	

<sup>36</sup> <https://consultations.ons.gov.uk/health-and-life-events/avoidable-mortality-definition/>

<sup>37</sup> <http://www.oecd.org/health/health-systems/Avoidable-mortality-2019-Joint-OECD-Eurostat-List-preventable-treatable-causes-of-death.pdf>

Condition Group & Cause	ICD-10 Codes	Age	Treatable	Preventable
<b>Neoplasms</b>				
Lip, oral cavity and pharynx cancer	C00-C14	0-74		•
Oesophageal cancer	C15	0-74		•
Stomach cancer	C16	0-74		•
Liver cancer	C22	0-74		•
Lung cancer	C33-C34	0-74		•
Mesothelioma	C45	0-74		•
Skin (melanoma) cancer	C43	0-74		•
Bladder cancer	C67	0-74		•
Cervical cancer	C53	0-74	• (50%)	• (50%)
Colorectal cancer	C18-C21	0-74	•	
Breast cancer (Female only)	C50	0-74	•	
Uterus cancer	C54,C55	0-74	•	
Testicular cancer	C62	0-74	•	
Thyroid cancer	C73	0-74	•	
Hodgkin's disease	C81	0-74	•	
Lymphoid leukaemia	C91.0, C91.1	0-74	•	
Benign neoplasm	D10-D36	0-74	•	
<b>Endocrine and metabolic diseases</b>				
Nutritional deficiency anaemia	D50-D53			•
Diabetes mellitus	E10-E14		• (50%)	• (50%)
Thyroid disorders	E00-E07		•	
Adrenal disorders	E24-E25 (except E24.4), E27		•	
<b>Diseases of the nervous system</b>				
Epilepsy	G40,G41	0-74	•	
<b>Diseases of the circulatory system</b>				
Aortic aneurysm	I71	0-74	• (50%)	• (50%)
Hypertensive diseases	I10-I13, I15	0-74	• (50%)	• (50%)
Ischaemic heart diseases	I20-I25	0-74	• (50%)	• (50%)
Cerebrovascular diseases	I60-I69	0-74	• (50%)	• (50%)
Other atherosclerosis	I70, I73.9	0-74	• (50%)	• (50%)
Rheumatic and other heart diseases	I00-I09	0-74	•	
Venous thromboembolism	I26, I80	0-74	•	

Condition Group & Cause	ICD-10 Codes	Age	Treatable	Preventable
<b>Diseases of the respiratory system</b>				
Influenza	J09-J11	0-74		•
Pneumonia due to Streptococcus pneumoniae or Haemophilus influenzae	J13-J14	0-74		•
Chronic lower respiratory diseases	J40-J44	0-74		•
Lung diseases due to external agents	J60-J64, J66-J70, J82, J92	0-74		•
Upper respiratory infections	J00-J06, J30-J39	0-74	•	
Pneumonia, not elsewhere classified or organism unspecified	J12, J15, J16- J18	0-74	•	
Acute lower respiratory infections	J20-J22	0-74	•	
Asthma and bronchiectasis	J45-J47	0-74	•	
Adult respiratory distress syndrome	J80	0-74	•	
Pulmonary oedema	J81	0-74	•	
Abscess of lung and mediastinum pyothorax	J85, J86	0-74	•	
Other pleural disorders	J90, J93, J94	0-74	•	
<b>Diseases of the digestive system</b>				
Gastric and duodenal ulcer	K25-K28	0-74	•	
Appendicitis	K35-K38	0-74	•	
Abdominal hernia	K40-K46	0-74	•	
Cholelithiasis and cholecystitis	K80-K81	0-74	•	
Other diseases of gallbladder or biliary tract	K82-K83	0-74	•	
Acute pancreatitis	K85.0, K85.1, K85.3, K85.8, K85.9	0-74	•	
Other diseases of pancreas	K86.1, K86.2, K86.3, K86.8, K86.9	0-74	•	
<b>Diseases of the genitourinary system</b>				
Nephritis and nephrosis	N00-N07	0-74	•	
Obstructive uropathy	N13, N20-N21, N35	0-74	•	
Renal failure	N17-N19	0-74	•	
Renal colic	N23	0-74	•	
Disorders resulting from renal tubular dysfunction	N25	0-74	•	
Unspecified contracted kidney, small kidney of unknown cause	N26-N27	0-74	•	
Inflammatory diseases of genitourinary system	N34.1, N70-N73, N75.0, N75.1, N76.4, N76.6	0-74	•	
Prostatic hyperplasia	N40	0-74	•	

Condition Group & Cause	ICD-10 Codes	Age	Treatable	Preventable
<b>Pregnancy, childbirth and perinatal period</b>				
Tetanus neonatorum	A33	0-74		•
Obstetrical tetanus	A34	0-74		•
Pregnancy, childbirth and the puerperium	O00-O99	0-74	•	
Certain conditions originating in the perinatal period	P00-P96	0-74	•	
<b>Congenital malformations</b>				
Certain congenital malformations (neural tube defects)	Q00, Q01, Q05	0-74		•
Congenital malformations of the circulatory system (heart defects)	Q20-Q28	0-74	•	
<b>Adverse effects of medical and surgical care</b>				
Drugs, medicaments and biological substances causing adverse effects in therapeutic use	Y40-Y59	0-74	•	
Misadventures to patients during surgical and medical care	Y60-Y69, Y83-Y84	0-74	•	
Medical devices associated with adverse incidents in diagnostic and therapeutic use	Y70-Y82	0-74	•	
<b>Injuries</b>				
Transport Accidents	V01-V99	0-74		•
Accidental Injuries	W00-X39, X46-59	0-74		•
Intentional self-harm	X66-X84	0-74		•
Event of undetermined intent	Y16-Y34	0-74		•
Assault	X86-Y09	0-74		•
<b>Alcohol related and drug-related deaths</b>				
Alcohol specific disorders and poisonings	E24.4, F10, G31.2, G62.1, G72.1, I42.6, K29.2, K70, K85.2, K86.0, Q86.0, R78.0, X45, X65, Y15	0-74		•
Other alcohol related disorders	K73, K74.0-K74.2, K74.6, K74.9	0-74		•
Drug disorders and poisonings	F11-F16, F18-F19, X40-X44, X85, Y10-Y14	0-74		•
Intentional self-poisoning by drugs	X60-X64	0-74		•

**Table 8: Admissions – Alcohol Related Causes<sup>38</sup>**

ICD-10 code	Description
E24.4	Alcohol induced Pseudo-Cushing's syndrome
E51.2	Wernicke's Encephalopathy
F10	Mental and Behavioural disorders due to use of alcohol
G31.2	Degeneration of the nervous system due to alcohol
G62.1	Alcoholic polyneuropathy
G72.1	Alcoholic myopathy
I42.6	Alcoholic cardiomyopathy
K29.2	Alcoholic gastritis
K70	Alcoholic liver disease
K85.2	Alcohol-induced acute pancreatitis
K86.0	Alcohol-induced chronic pancreatitis
O35.4	Maternal care for (suspected) damage to foetus from alcohol
P04.3	Foetus and newborn affected by maternal use of alcohol
Q86.0	Foetal alcohol syndrome (dysmorphic)
T51.0	Toxic effect of ethanol
T51.1	Toxic effect of methanol
T51.9	Toxic effect of alcohol, unspecified
X45	Accidental poisoning by and exposure to alcohol
X65	Intentional self-poisoning by and exposure to alcohol
Y15	Poisoning by and exposure to alcohol, undetermined intent
Y57.3	Alcohol deterrents
Y90	Evidence of alcohol involvement determined by blood alcohol level
Y91	Evidence of alcohol involvement determined by level intoxication
Z50.2	Alcohol rehabilitation
Z71.4	Alcohol abuse counselling and surveillance
Z72.1	Alcohol use

<sup>38</sup> The definition for admissions due to alcohol related causes was updated for 2017/18 to include ICD Code K85.2: alcohol-induced chronic pancreatitis.

**Table 9: Deaths – Alcohol Specific Causes**

ICD-10 code	Description
E24.4	Alcohol-induced pseudo-Cushing's syndrome
F10	Mental and Behavioural disorders due to use of alcohol
G31.2	Degeneration of the nervous system due to alcohol
G62.1	Alcoholic polyneuropathy
G72.1	Alcohol myopathy
I42.6	Alcoholic cardiomyopathy
K29.2	Alcoholic gastritis
K70	Alcoholic liver disease
K85.2	Alcohol-induced acute pancreatitis
K86.0	Alcohol induced chronic pancreatitis
Q86.0	Fetal alcohol syndrome (dysmorphic)
R78.0	Excess alcohol blood levels
X45	Accidental poisoning by and exposure to alcohol
X65	Intentional self-poisoning by and exposure to alcohol
Y15	Poisoning by and exposure to alcohol, undetermined intent

**Table 10: Admissions and Deaths – Drug Related Causes**

ICD-10 code	Description
F11-16, F18-F19	Mental and Behavioural disorders due to drug use (excluding alcohol and tobacco)
X40-44	Accidental poisoning by drugs, medicaments and biological substances
X60-X64	Intentional self-poisoning by drugs, medicaments and biological substances
X85	Assault by drugs, medicaments and biological substances
Y10-14	Poisoning by drugs, medicaments and biological substances, undetermined intent

**Table 11: Deaths – Drugs Misuse**

ICD-10 code	Description
F11-16, F19	Mental and Behavioural disorders due to drug use (excluding alcohol, tobacco and volatile substances)
Deaths due to the following categories <i>and</i> where a drug controlled under the Misuse of Drugs Act 1971 was mentioned;	
X40-44	Accidental poisoning by drugs, medicaments and biological substances
X60-X64	Intentional self-poisoning by drugs, medicaments and biological substances
X85	Assault by drugs, medicaments and biological substances
Y10-14	Poisoning by drugs, medicaments and biological substances, undetermined intent

Table 12: Deaths – Smoking Related Causes<sup>39</sup>

Cause of Death	ICD-10 code	Attributable Percentage	
		Men	Women
<b>Cancer</b>			
Lung	C33 – C34	90%	79%
Upper respiratory	C32, C14.0	77%	58%
Oesophagus	C15	70%	72%
Bladder	C67	49%	20%
Kidney	C64	41%	7%
Stomach	C16	35%	10%
Pancreas	C25	26%	30%
Unspecified site	C80	33%	7%
Myeloid Leukaemia	C92	19%	10%
<b>Respiratory</b>			
Chronic obstructive lung disease	J44	87%	83%
Pneumonia 35-64	J18	33%	53%
Pneumonia 65+	J18	23%	13%
<b>Circulatory</b>			
Ischaemic heart disease 35-54	I20-I25	55%	63%
Ischaemic heart disease 55-64	I20-I25	41%	36%
Ischaemic heart disease 65-74	I20-I25	25%	18%
Ischaemic heart disease 75+	I20-I25	9%	5%
Cerebrovascular disease 35-54	I60-I69	56%	53%
Cerebrovascular disease 55-64	I60-I69	33%	38%
Cerebrovascular disease 65-74	I60-I69	16%	31%
Cerebrovascular disease 75+	I60-I69	4%	2%
Aortic Aneurysm	I71	64%	66%
Myocardial Degeneration	I51.5	27%	18%
Atherosclerosis	I70	21%	21%
<b>Digestive</b>			
Stomach/Duodenum Ulcer	K25-K26	53%	59%
<b>Disease Prevented by Smoking</b>			
Parkinson's Disease	G20	-51%	-30%
Endometrial Cancer	C54	N/A	-16%

<sup>39</sup>It should be noted that this definition is specific to the death rates in NI and therefore differs from those used in other parts of the UK and other countries, meaning it is not directly comparable.

The table below lists the previous definition for all causes of death considered avoidable, with indication as to which are considered amenable, preventable or both. This list is for use with the additional indicators in [Appendix C](#) which include the avoidable mortality indicator as used by the NI Programme for Government.

**Table 13: Amenable, Preventable and Avoidable Mortality (Previous ONS Definition).**

Condition Group & Cause	ICD-10 Codes	Age	Amenable	Preventable
<b>Infections</b>				
Tuberculosis	A15-A19, B90	0-74	•	•
Selected invasive bacterial and protozoal infections	A38-A41, A46, A48.1, B50-B54, G00, G03, J02, L03	0-74	•	
Hepatitis C	B17.1, B18.2	0-74	•	•
Pertussis (whooping cough)	A37	0-14	•	•
Measles	B05	1-14	•	•
Rubella	B06	0-14		•
Other infections (Diphtheria, Tetanus, Poliomyelitis and Varicella)	A35, A36, A80, B01	0-19	•	•
Intestinal infections	A00-A09	0-14	•	
HIV/AIDS	B20-B24	All	•	•
<b>Neoplasms</b>				
Malignant neoplasm of lip, oral cavity and pharynx	C00-C14	0-74		•
Malignant neoplasm of oesophagus	C15	0-74		•
Malignant neoplasm of stomach	C16	0-74		•
Malignant neoplasm of colon and rectum	C18-C21	0-74	•	•
Malignant neoplasm of liver	C22	0-74		•
Malignant neoplasm of trachea, bronchus and lung	C33-C34	0-74		•
Malignant melanoma of skin	C43	0-74	•	•
Mesothelioma	C45	0-74		•
Malignant neoplasm of breast	C50	0-74	•	•
Malignant neoplasm of cervix uteri	C53	0-74	•	•
Malignant neoplasm of bladder	C67	0-74	•	
Malignant neoplasm of thyroid gland	C73	0-74	•	
Hodgkin's disease	C81	0-74	•	
Leukaemia	C91, C92.0	0-44	•	
Malignant neoplasm of testis	C62	0-74	•	
Malignant neoplasm of unspecified parts of uterus and body of uterus	C54-C55	0-44	•	
Benign neoplasms	D10-D36	0-74	•	
<b>Nutritional, endocrine and metabolic</b>				
Diabetes mellitus	E10-E14	0-74	•	•
Diseases of Thyroid	E00-E07	0-74	•	
Addison's Disease	E27.1	0-74	•	



Condition Group & Cause	ICD-10 Codes	Age	Amenable	Preventable
<b>Drug use disorders</b>				
Alcohol related diseases, excluding external causes	F10, G31.2, G62.1, I42.6, K29.2, K70, K73, K74 (excl. K74.3-K74.5), K86.0	0-74		•
Illicit drug use disorders	F11-F16, F18-F19	0-74		•
<b>Neurological disorders</b>				
Epilepsy and status epilepticus	G40-G41	0-74	•	
<b>Cardiovascular diseases</b>				
Rheumatic and other valvular heart disease	I01-I09	0-74	•	
Hypertensive diseases	I10-I15	0-74	•	
Ischaemic heart disease	I20-I25	0-74	•	•
DVT with pulmonary embolism	I26, I80.1-I80.3, I80.9, I82.9	0-74		•
Cerebrovascular diseases	I60-I69	0-74	•	
Aortic aneurysm and dissection	I71	0-74		•
<b>Respiratory diseases</b>				
Influenza (including swine flu)	J09-J11	0-74	•	•
Pneumonia	J12-J18	0-74	•	
Chronic obstructive pulmonary disorder	J40-J44	0-74	•	•
Asthma	J45-J46	0-74	•	
Selected respiratory diseases	J00-J06, J20-J22, J30-J39	1-14	•	
<b>Digestive disorders</b>				
Gastric and duodenal ulcer	K25-K28	0-74	•	
Acute abdomen, appendicitis, intestinal obstruction, cholecystitis/lithiasis, pancreatitis, hernia	K35-K38, K40-K46, K80-K83, K85, K86.1-K86.9, K91.5	0-74	•	
<b>Genitourinary disorders</b>				
Nephritis and nephrosis	N00-N07, N17-N19, N25-N27	0-74	•	
Obstructive uropathy and prostatic hyperplasia	N13, N20-N21, N35, N40, N99.1	0-74	•	
<b>Maternal and infant</b>				
Complications of perinatal period	P00-P96, A33	All	•	
Congenital malformations of the circulatory system	Q20-Q28	0-74	•	
Spina Bifida	Q05	0-74		•
<b>Unintentional injuries</b>				
Transport Accidents	V01-V99	All		•
Accidental Injury	W00-X59	All		•
<b>Intentional injuries</b>				
Suicide and self inflicted injuries	X60-X84, Y10-Y34	All		•
Homicide/Assault	X85-Y09, U50.9	All		•
Misadventures to patients during surgical and medical care	Y60-Y69, Y83-Y84	All	•	•

## **Also available for the Health & Social Care Inequalities Monitoring System (HSCIMS)**

### **Life Expectancy in Northern Ireland**

Presenting the latest official estimates of life expectancy in Northern Ireland and an examination of the causes that contribute to the change in life expectancy over time, as well as the differentials between gender and deprivation classification and across HSC Trusts and Local Government Districts. The latest figures for life expectancy at 65, healthy life expectancy and disability-free life expectancy are also included.

<https://www.health-ni.gov.uk/articles/life-expectancy-northern-ireland>

### **Making Life Better**

#### **Monitoring the Wider Social Determinants of Health & Wellbeing Key Indicators**

Monitoring of the key indicators of the wider social determinant of health & wellbeing set out against each of the themes contained in the making life better strategic framework (Annual).

<http://www.health-ni.gov.uk/articles/social-determinants-health-statistics>

### **Coronavirus Related Health Inequalities**

Presenting an analysis of coronavirus (Covid-19) related health inequalities by assessing differences in positive COVID-19 test cases, COVID-19 admissions and deaths between the most and least deprived areas of Northern Ireland. An assessment of variations across age, sex and urban & rural areas is also included.

<https://www.health-ni.gov.uk/articles/coronavirus-related-health-inequalities>

#### **Produced by:**

Public Health Information & Research Branch, Information Analysis Directorate

Department of Health

Castle Buildings, Belfast BT4 3SQ

Tel: 02890522501

Email: [healthinequalities@health-ni.gov.uk](mailto:healthinequalities@health-ni.gov.uk)

<https://www.health-ni.gov.uk/articles/health-inequalities-statistics>